

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 39/12, 37/02, C12N 15/00, C07K 3/00, C07H 15/12		A1	(11) International Publication Number: WO 94/28929
		(43) International Publication Date: 22 December 1994 (22.12.94)	
(21) International Application Number: PCT/US94/06036		(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KG, KP, KR, KZ, LK, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 7 June 1994 (07.06.94)			
(30) Priority Data: 072,833 7 June 1993 (07.06.93) US			
(60) Parent Application or Grant (63) Related by Continuation US 072,833 (CIP) Filed on 7 June 1993 (07.06.93)		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(71) Applicant (for all designated States except US): GENENTECH, INC. [US/US]; 460 Point San Bruno Boulevard, South San Francisco, CA 94080 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): BERMAN, Phillip, W. [US/US]; 95 Cheyenne Point, Portola Valley, CA 94028 (US). NAKAMURA, Gerald, R. [US/US]; 1923 29th Avenue, San Francisco, CA 94116 (US).			
(74) Agents: TERLIZZI, Laura et al.; Skjerven, Morrill, Macpherson, Franklin & Friel, Suite 700, 25 Metro Drive, San Jose, CA 95110 (US).			

(54) Title: HIV ENVELOPE POLYPEPTIDES

(57) Abstract

A method for the rational design and preparation of vaccines based on HIV envelope polypeptides is described. In one embodiment, the method for making an HIV gp120 subunit vaccine for a geographic region comprises determining neutralizing epitopes in the V2 and/or C4 domains of gp120 of HIV as depicted in the figure. In a preferred embodiment of the method, neutralizing epitopes for the V2, V3 and C4 domains of gp120 are determined. Also described are DNA sequences encoding gp120 from preferred vaccine strains of HIV.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

HIV ENVELOPE POLYPEPTIDES

FIELD OF THE INVENTION

5 This invention relates to the rational design and
preparation of HIV vaccines based on HIV envelope
polypeptides and the resultant vaccines. This
invention further relates to improved methods for HIV
serotyping and immunogens which induce antibodies
10 useful in the serotyping methods.

BACKGROUND OF THE INVENTION

Acquired immunodeficiency syndrome (AIDS) is
caused by a retrovirus identified as the human
15 immunodeficiency virus (HIV). There have been intense
effort to develop a vaccine. These efforts have
focused on inducing antibodies to the HIV envelope
protein. Recent efforts have used subunit vaccines
where an HIV protein, rather than attenuated or killed
20 virus, is used as the immunogen in the vaccine for
safety reasons. Subunit vaccines generally include
gp120, the portion of the HIV envelope protein which is
on the surface of the virus.

The HIV envelope protein has been extensively
25 described, and the amino acid and RNA sequences
encoding HIV envelope from a number of HIV strains are
known (Myers, G. et al., 1992. Human Retroviruses and
AIDS. A compilation and analysis of nucleic acid and
amino acid sequences. Los Alamos National Laboratory,
30 Los Alamos, New Mexico). The HIV envelope protein is a
glycoprotein of about 160 kd (gp160) which is anchored
in the membrane bilayer at its carboxyl terminal
region. The N-terminal segment, gp120, protrudes into
the aqueous environment surrounding the virion and the
35 C-terminal segment, gp41, spans the membrane. Via a
host-cell mediated process, gp160 is cleaved to form

gp120 and the integral membrane protein gp41. As there is no covalent attachment between gp120 and gp41, free gp120 is released from the surface of virions and infected cells.

5 The gp120 molecule consists of a polypeptide core of 60,000 daltons which is extensively modified by N-linked glycosylation to increase the apparent molecular weight of the molecule to 120,000 daltons. The amino acid sequence of gp120 contains five
10 relatively conserved domains interspersed with five hypervariable domains. The positions of the 18 cysteine residues in the gp120 primary sequence, and the positions of 13 of the approximately 24 N-linked glycosylation sites in the gp120 sequence are common to
15 all gp120 sequences. The hypervariable domains contain extensive amino acid substitutions, insertions and deletions. Sequence variations in these domains result in up to 30% overall sequence variability between gp120 molecules from the various viral isolates. Despite
20 this variation, all gp120 sequences preserve the virus's ability to bind to the viral receptor CD4 and to interact with gp41 to induce fusion of the viral and host cell membranes.

gp120 has been the object of intensive
25 investigation as a vaccine candidate for subunit vaccines, as the viral protein which is most likely to be accessible to immune attack. gp120 is considered to be a good candidate for a subunit vaccine, because (i) gp120 is known to possess the CD4 binding domain by
30 which HIV attaches to its target cells, (ii) HIV infectivity can be neutralized *in vitro* by antibodies to gp 120, (iii) the majority of the *in vitro* neutralizing activity present in the serum of HIV infected individuals can be removed with a gp120
35 affinity column, and (iv) the gp120/gp41 complex

appears to be essential for the transmission of HIV by cell-to-cell fusion.

The identification of epitopes recognized by virus neutralizing antibodies is critical for the rational design of vaccines effective against HIV-1 infection. One way in which antibodies would be expected to neutralize HIV-1 infection is by blocking the binding of the HIV-1 envelope glycoprotein, gp120, to its cellular receptor, CD4. However, it has been surprising that the CD4 blocking activity, readily demonstrated in sera from HIV-1 infected individuals (31, 44) and animals immunized with recombinant envelope glycoproteins (1-3), has not always correlated with neutralizing activity (2, 31, 44). Results obtained with monoclonal antibodies have shown that while some of the monoclonal antibodies that block the binding of gp120 to CD4 possess neutralizing activity, others do not (4, 7, 16, 26, 33, 35, 43, 45). When the neutralizing activity of CD4 blocking monoclonal antibodies are compared to those directed to the principal neutralizing determinant (PND) located in the third variable domain (V3 domain) of gp120 (10, 39), the CD4 blocking antibodies appear to be significantly less potent. Thus, CD4 blocking monoclonal antibodies typically exhibit 50% inhibitory concentration values (IC_{50}) in the 1-10 μ g/ml range (4, 16, 26, 33, 35, 43, 45) whereas PND directed monoclonal antibodies typically exhibit IC_{50} values in the 0.1 to 1.0 μ g/ml range (23, 33, 42).

Subunit vaccines, based on gp120 or another viral protein, that can effectively induce antibodies that neutralize HIV are still being sought. However, to date no vaccine has not been effective in conferring protection against HIV infection.

35

DESCRIPTION OF THE BACKGROUND ART

Recombinant subunit vaccines are described in Berman et al., PCT/US91/02250 (published as number WO91/15238 on 17 October 1991). See also, e.g. Hu et al., *Nature* 328:721-724 (1987) (vaccinia virus-HIV envelope recombinant vaccine); Arthur et al., *J. Virol.* 63(12): 5046-5053 (1989) (purified gp120); and Berman et al., *Proc. Natl. Acad. Sci. USA* 85:5200-5204 (1988) (recombinant envelope glycoprotein gp120).

10 Numerous sequences for gp120 are known. The sequence of gp120 from the IIIB substrain of HIV-1_{LAI} referred to herein is that determined by Muesing et al., "Nucleic acid structure and expression of the human AIDS/lymphadenopathy retrovirus, *Nature* 313:450-458 (1985). The sequences of gp120 from the NY-5, JrCSF, Z6, Z321, and HXB2 strains of HIV-1 are listed by Myers et al., "Human Retroviruses and AIDS; A compilation and analysis of nucleic acid and amino acid sequences," Los Alamos National Laboratory, Los Alamos, New Mexico (1992). The sequence of the Thai isolate A244 is provided by McCutchan et al., "Genetic Variants of HIV-1 in Thailand," *AIDS Res. and Human Retroviruses* 8:1887-1895 (1992). The MN₁₉₈₄ clone is described by Gargo et al., "Envelope sequences of two new United States HIV-1 isolates," *Virol.* 164: 531-536 (1988). The amino acid sequence of this MN clone differs by approximately 2% from the MN-gp120 clone (MN_{ONE}) disclosed herein and obtained by Berman et al.

15
20
25
30

Each of the above-described references is incorporated herein by reference in its entirety.

SUMMARY OF THE INVENTION

The present invention provides a method for the rational design and preparation of vaccines based on HIV envelope polypeptides. This invention is based on the discovery that there are neutralizing epitopes in

35

the V2 and C4 domains of gp120, in addition to the neutralizing epitopes in the V3 domain. In addition, the amount of variation of the neutralizing epitopes is highly constrained, facilitating the design of an HIV subunit vaccine that can induce antibodies that neutralize a plurality of HIV strains for a given geographic region.

In one embodiment, the present invention provides a method for making an HIV gp120 subunit vaccine for a geographic region in which a neutralizing epitope in the V2 and/or C4 domains of gp120 of HIV isolates from the geographic region is determined and an HIV strain having gp120 which has a neutralizing epitope in the V2 or C4 domain which is common among isolates in the geographic region is selected and used to make the vaccine.

In a preferred embodiment of the method, neutralizing epitopes for the V2, V3, and C4 domains of gp120 from HIV isolates from the geographic region are determined. At least two HIV isolates having different neutralizing epitopes in the V2, V3, or C4 domain are selected and used to make the HIV gp120 subunit vaccine. Preferably, each of the selected isolates have one of the most common neutralizing epitopes for the V2, V3, or C4 domains.

The invention also provides a multivalent HIV gp120 subunit vaccine. The vaccine comprises gp120 from two isolates of HIV having at least one different neutralizing epitope. Preferably, the isolates have the most common neutralizing epitopes in the geographic region for one of the domains.

A DNA sequence of less than 5 kilobases encoding gp120 from preferred vaccine strains of HIV, GNE₁ and GNE₁₆, expression construct comprising the GNE₁-gp120 and GNE₁₆-gp120 encoding DNA under the transcriptional and translational control of a heterologous promoter, and

isolated GNE₈-gp120 and GNE₁₆-gp120 are also provided. The invention further provides improved methods for HIV serotyping in which epitopes in the V2 or C4 domains of gp120 are determined and provides immunogens (truncated
5 gp120 sequences) which induce antibodies useful in the serotyping methods.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1 describes inhibition of CD4 binding by
10 monoclonal antibodies to recombinantly produced gp120 from the MN strain of HIV (MN-rgp120). Mice were immunized with MN-rgp120 and the resulting splenocytes were fused with the NP3X63.Ag8.653 cell line as described in Example 1. Thirty-five stable hybridoma
15 clones, reactive with MN-rgp120 were identified by ELISA. Secondary screening revealed seven cell lines (1024, 1093, 1096, 1097, 1110, 1112, and 1027) secreting antibodies able to inhibit the binding of MN-rgp120 to biotin labeled recombinantly produced CD4
20 (rsCD4) in a ELISA using HRPO-strepavidin. Data obtained with monoclonal antibodies from the same fusion (1026, 1092, 1126) that failed to inhibit MN-rgp120 binding to CD4 is shown for purposes of comparison.

25 FIGURE 2 shows neutralizing activity of CD4-blocking monoclonal antibodies to MN-rgp120. Monoclonal antibodies that blocked the binding of MN-rgp120 to CD4 were screened for the capacity to inhibit the infection of MT2 cells by the MN strain of
30 HIV-1 in vitro. Cell free virus was added to wells containing serially diluted antibodies and incubated at 4°C for 1 hr. After incubation, MT-2 cells were added to the wells and the cultures were then grown for 5 days at 37°C. Cell viability was then measured by
35 addition of the colorimetric tetrazolium compound MTT as described in reference (35) of Example 1. The

optical densities of each well were measured at 540 nm using a microtiter plate reading spectrophotometer. Inhibition of virus infectivity was calculated by dividing the mean optical densities from wells containing monoclonal antibodies by the mean value of wells that received virus alone. Monoclonal antibodies that blocked CD4 binding are the same as those indicated in Figure Legend 1. Data from the V3-directed monoclonal antibody to MN-rgp120 (1034) is provided as a positive control. Data obtained with the V3 directed monoclonal antibody, 11G5, specific for the IIIB strain of HIV-1 (33) is shown as a negative control.

FIGURE 3 is a diagram of gp120 fragments used to localize the epitopes recognized by the CD4 blocking monoclonal antibodies to MN-rgp120. A series of fragments (A) corresponding to the V4 and C4 domains (B) (SEQ. ID. NO. 14) of the gene encoding MN-rgp120 were prepared by PCR. The gp120 gene fragments were fused to a fragment of the gene encoding Herpes Simplex Virus Type 1 glycoprotein D that encoded the signal sequence and 25 amino acids from the mature amino terminus. The chimeric genes were assembled into a mammalian cell expression vector (PRK5) that provided a CMV promoter, translational stop codons and an SV40 polyadenylation site. The embryonic human kidney adenocarcinoma cell line, 293s, was transfected with the resulting plasmid and recombinant proteins were recovered from growth conditioned cell culture medium. Fragments of MN-rgp120, expressed as HSV-1 Gd fusion proteins, were produced by transient transfection of 293s cells (Example 1). To verify expression, cells were metabolically labeled with [³⁵S]-methionine, and the resulting growth conditioned cell culture supernatants were immunoprecipitated (C) using a monoclonal antibody, 5B6, specific for the amino

terminus of HSV-1 Gd and fixed *S. aureus*. The immunoprecipitated proteins were resolved on 4 to 20 % acrylamide gradient gels using SDS-PAGE and visualized by autoradiography. The samples were: Lane 1, 5 FMN.368-408; lane 2, FMN.368-451; lane 3, FMN.419-443; lane 4, FMN.414-451; lane 5, MN-rgp120. The gel demonstrated that the proteins were expressed and migrated at the expected molecular weights.

FIGURE 4 shows a C4 domain sequence comparison 10 (SEQ. ID. Nos. 3-13). The C4 domain amino acid sequences of recombinant and virus derived gp120s used for monoclonal antibody binding studies were aligned starting the amino terminal cysteine. Amino acid positions are designated with respect to the sequence 15 of MN-rgp120. Sequences of the LAI substrains, IIIB, BH10, Bru, HXB2, and HXB3 are shown for purposes of comparison.

FIGURE 5 shows sequences of C4 domain mutants of MN-rgp120 (SEQ. ID. Nos. 3 and 15-23). Nucleotide 20 substitutions, resulting in the amino acid sequences indicated, were introduced into the C4 domain of MN-rgp120 gene using recombinant PCR. The resulting variants were assembled into the expression plasmid, pRK5, which was then transfected into 293s cells. The 25 binding of monoclonal antibodies to the resulting C4 domain variants was then analyzed (Table 5) by ELISA.

FIGURE 6 illustrates the reactivity of monoclonal antibody 1024 with HIV-1_{LAI} substrains. The cell surface binding of the C4 domain reactive monoclonal 30 antibody 1024 to H9 cells chronically infected with the IIIB, HXB2, HXB3, and HXB10 substrains of HIV-1 LAI or HIV-1MN was analyzed by flow cytometry. Cultures of virus infected cells were reacted with either monoclonal antibody 1024, a nonrelevant monoclonal 35 antibody (control), or a broadly cross reactive monoclonal antibody (1026) raised against rgp120.

After washing away unbound monoclonal antibody, the cells were then labeled with fluorescein conjugated goat antibody to mouse IgG (Fab'), washed and fixed with paraformaldehyde. The resulting cells were
5 analyzed for degree of fluorescence intensity using a FACSCAN (Becton Dickenson, Fullerton, CA). Fluorescence was measured as mean intensity of the cells expressed as mean channel number plotted on a log scale.

10 FIGURE 7 shows the determination of the binding affinity of monoclonal antibodies for MN-rgp120. CD4 blocking monoclonal antibodies raised against MN-rgp120 (1024 and 1097) or IIIB-rgp120 (13H8 and 5C2) were labeled with [¹²⁵I] and binding titrations using
15 MN-rgp120 (A and B) or IIIB-rgp120 (C and D) were carried out as described in the Example 1. A, binding of monoclonal antibody 1024; B binding of monoclonal antibody 1097; C, binding of monoclonal antibody 13H8; and D binding of monoclonal antibody 5C2.

20 FIGURE 8 shows the correlation between gp120 binding affinity (K_d) and neutralizing activity (IC_{50}) of monoclonal antibodies to the C4 domain of MN-rgp120. Binding affinities of monoclonal antibodies to the C4 domain of gp120 were determined by Scatchard analysis
25 (Figure 9, Table 5). The resulting values were plotted as a function of the log of their neutralizing activities (IC_{50}) determined in Figure 2 and Table 6.

FIGURE 9 depicts the amino acid sequence of the mature envelope glycoprotein (gp120) from the MN_{GNE}
30 clone of the MN strain of HIV-1 (SEQ. ID. NO. 1). Hypervariable domains are from 1-29 (signal sequence), 131-156, 166-200, 305-332, 399-413, and 460-469. The V and C regions are indicated (according to Modrow et al., *J. Virology* 61(2):570 (1987). Potential
35 glycosylation sites are marked with a (*).

FIGURE 10 depicts the amino acid sequence of a fusion protein of the residues 41-511 of the mature envelope glycoprotein (gp120) from the MN_{ONE} clone of the MN strain of HIV-1, and the gD-1 amino terminus from the herpes simplex glycoprotein gD-1. (SEQ. ID. NO. 2). The V and C regions are indicated (according to Modrow et al., *J. Virology* 61(2):570 (1987)). Potential glycosylation sites are marked with a (*).

10 DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a method for the rational design and preparation of vaccines based on HIV envelope polypeptides. This invention is based on the discovery that there are neutralizing epitopes in the V2 and C4 domains of gp120, in addition to the neutralizing epitopes in the V3 domain. Although the amino acid sequences of the neutralizing epitopes in the V2, V3, and C4 domains are variable, it has now been found that the amount of variation is highly constrained. The limited amount of variation facilitates the design of an HIV subunit vaccine that can induce antibodies that neutralize the most common HIV strains for a given geographic region. In particular, the amino acid sequence of neutralizing epitopes in the V2, V3, and C4 domains for isolates of a selected geographic region is determined. gp120 from isolates having the most common neutralizing epitope sequences are utilized in the vaccine.

The invention also provides a multivalent gp120 subunit vaccine wherein gp120 present in the vaccine is from at least two HIV isolates which have different amino acid sequences for a neutralizing epitope in the V2, V3, or C4 domain of gp120. The invention further provides improved methods for HIV serotyping in which epitopes in the V2 or C4 domains of gp120 are

determined and provides immunogens which induce antibodies useful in the s rotyping methods.

The term "subunit vaccine" is used herein, as in the art, to refer to a viral vaccine that does not contain virus, but rather contains one or more viral proteins or fragments of viral proteins. As used herein, the term "multivalent" means that the vaccine contains gp120 from at least two HIV isolates having different amino acid sequences for a neutralizing epitope.

Vaccine Design Method

The vaccine design method of this invention is based on the discovery that there are neutralizing epitopes in the V2 and C4 domains of gp120, in addition to those found in the principal neutralizing domain (PND) in the V3 domain. Selecting an HIV isolate with appropriate neutralizing epitopes in the V2 and/or C4 domains provides a vaccine that is designed to induce immunity to the HIV isolates present in a selected geographic region. In addition, although the amino acid sequence of the V2, V3, and C4 domains containing the neutralizing epitopes is variable, the amount of variation is highly constrained, facilitating the design of a multivalent vaccine which can neutralize a plurality of the most common HIV strains for a given geographic region.

The method for making an HIV gp120 subunit vaccine depends on the use of appropriate strains of HIV for a selected geographic region. Appropriate strains of HIV for the region are selected by determining the neutralizing epitopes for HIV isolates and the percentage of HIV infections attributable to each strain present in the region. HIV strains which have the most common neutralizing epitopes in the V2 or C4 domains in the geographic region are selected.

Preferably, isolates that confer protection against the most common neutralizing epitopes in the V2, V3, and C4 domains for a geographic region are selected.

One embodiment of the method for making an HIV gp120 subunit vaccine from appropriate strains of HIV for a geographic region comprises the following steps. A neutralizing epitope in the V2 or C4 domain of gp120 of HIV isolates from the geographic region is determined. An HIV strain having gp120 with a neutralizing epitope in the V2 or C4 domain that is common among HIV isolates in the geographic region is selected. gp120 from the selected isolate is used to make an HIV gp120 subunit vaccine.

In another embodiment of the method, the neutralizing epitopes in the V2, V3, and C4 domains of gp120 from HIV isolates from the geographic region are determined. At least two HIV isolates having different neutralizing epitopes in the V2, V3, or C4 domain are selected and used to make an HIV gp120 subunit vaccine. Preferably, the vaccine contains gp120 from at least the two or three HIV strains having the most common neutralizing epitopes for the V2, V3, or C4 domains. More preferably, the vaccine contains gp120 from sufficient strains so that at least about 50%, preferably about 70%, more preferably about 80% or more of the neutralizing epitopes for the V2, V3, and C4 domains in the geographic region are included in the vaccine. The location of the neutralizing epitopes in the V3 region are well known. The location of the neutralizing epitopes in the V2 and C4 regions are described hereinafter.

Each of the steps of the method are described in detail below.

Determining neutralizing epitopes

The first step in designing a vaccine for a selected geographic region is to determine the neutralizing epitopes in the gp120 V2 and/or C4 domains. In a preferred embodiment, neutralizing epitopes in the V3 domain (the principal neutralizing domain) are also determined. The location of neutralizing epitopes in the V3 domain is well known. Neutralizing epitopes in the V2 and C4 domains have now been found to be located between about residues 163 and 200 and between about residues 420 and 440, respectively. In addition, the critical residues for antibody binding are residues 171, 173, 174, 177, 181, 183, 187, and 188 in the V2 domain and residues 429 and 432 in the C4 domain, as described in detail in the Examples.

The neutralizing epitopes for any isolate can be determined by sequencing the region of gp120 containing the neutralizing epitope. Alternatively, when antibodies specific for the neutralizing epitope, preferably monoclonal antibodies, are available the neutralizing epitope can be determined by serological methods as described hereinafter. A method for identification of additional neutralizing epitopes in gp120 is described hereinafter.

When discussing the amino acid sequences of various isolates and strains of HIV, the most common numbering system refers to the location of amino acids within the gp120 protein using the initiator methionine residue as position 1. The amino acid numbering reflects the mature HIV-1 gp120 amino acid sequence as shown by Figures 9 and Fig. 10 [SEQ. ID Nos. 1 and 2]. For gp120 sequences derived from other HIV isolates and which include their native HIV N-terminal signal sequence, numbering may differ. Although the nucleotide and amino acid residue numbers may not be applicable in other strains where upstream deletions or

insertions change the length of the viral genome and gp120, the region encoding the portions of gp120 is readily identified by reference to the teachings herein. The variable (V) domains and conserved (C) domains of gp120 are specified according to the nomenclature of Modrow et al. "Computer-assisted analysis of envelope protein sequences of seven human immunodeficiency virus isolates: predictions of antigenic epitopes in conserved and variable regions," *J. Virol.* 61:570-578 (1987).

The first step in identifying the neutralizing epitopes for any region of gp120 is to immunize an animal with gp120 to induce anti-gp120 antibodies. The antibodies can be polyclonal or, preferably, monoclonal. Polyclonal antibodies can be induced by administering to the host animal an immunogenic composition comprising gp120. Preparation of immunogenic compositions of a protein may vary depending on the host animal and the protein and is well known. For example, gp120 or an antigenic portion thereof can be conjugated to an immunogenic substance such as KLH or BSA or provided in an adjuvant or the like. The induced antibodies can be tested to determine whether the composition is specific for gp120. If a polyclonal antibody composition does not provide the desired specificity, the antibodies can be fractionated by ion exchange chromatography and immunoaffinity methods using intact gp120 or various fragments of gp120 to enhance specificity by a variety of conventional methods. For example, the composition can be fractionated to reduce binding to other substances by contacting the composition with gp120 affixed to a solid substrate. Those antibodies which bind to the substrate are retained. Fractionation techniques using antigens affixed to a variety of solid substrates such as affinity chromatography materials

including Sephadex, Sephar se and the like are well known.

Monoclonal anti-gp120 antibodies can be produced by a number of conventional methods. A mouse can be
5 injected with an immunogenic composition containing gp120 and spleen cells obtained. Those spleen cells can be fused with a fusion partner to prepare hybridomas. Antibodies secreted by the hybridomas can be screened to select a hybridoma wherein the
10 antibodies neutralize HIV infectivity, as described hereinafter. Hybridomas that produce antibodies of the desired specificity are cultured by standard techniques.

Infected human lymphocytes can be used to prepare
15 human hybridomas by a number of techniques such as fusion with a murine fusion partner or transformation with EBV. In addition, combinatorial libraries of human or mouse spleen can be expressed in E. coli to produce the antibodies. Kits for preparing
20 combinatorial libraries are commercially available. Hybridoma preparation techniques and culture methods are well known and constitute no part of the present invention. Exemplary preparations of monoclonal antibodies are described in the Examples.

25 Following preparation of anti-gp120 monoclonal antibodies, the antibodies are screened to determine those antibodies which are neutralizing antibodies. Assays to determine whether a monoclonal antibody neutralizes HIV infectivity are well known and are
30 described in the literature. Briefly, dilutions of antibody and HIV stock are combined and incubated for a time sufficient for antibody binding to the virus. Thereafter, cells that are susceptible to HIV infection are combined with the virus/antibody mixture and
35 cultured. MT-2 cells or H9 cells are susceptible to infection by m st HIV strains that are adapted for

growth in the laboratory. Activated peripheral blood mononuclear cells (PBMCs) or macrophages can be infected with primary isolates (isolates from a patient specimens which have not been cultured in T-cell lines or transformed cell lines). Daar et al, *Proc. Natl. Acad. Sci. USA* 87:6574-6578 (1990) describe methods for infecting cells with primary isolates.

After culturing the cells for about five days, the number of viable cells is determined, as by measuring metabolic conversion of the formazan MTT dye. The percentage of inhibition of infectivity is calculated to determine those antibodies that neutralize HIV. An exemplary preferred procedure for determining HIV neutralization is described in the Examples.

Those monoclonal antibodies which neutralize HIV are used to map the epitopes to which the antibodies bind. To determine the location of a gp120 neutralizing epitope, neutralizing antibodies are combined with fragments of gp120 to determine the fragments to which the antibodies bind. The gp120 fragments used to localize the neutralizing epitopes are preferably made by recombinant DNA methods as described hereinafter and exemplified in the Examples. By using a plurality of fragments, each encompassing different, overlapping portions of gp120, an amino acid sequence encompassing a neutralizing epitope to which a neutralizing antibody binds can be determined. A preferred exemplary determination of the neutralizing epitopes to which a series of neutralizing antibodies binds is described in detail in the Examples.

This use of overlapping fragments can narrow the location of the epitope to a region of about 20 to 40 residues. To confirm the location of the epitope and narrow the location to a region of about 5 to 10 residues, site-directed mutagenicity studies are preferably performed. Such studies can also determine

the critical residues for binding of neutralizing antibodies. A preferred exemplary site-directed mutagenicity procedure is described in the Examples.

To perform site-directed mutagenicity studies, recombinant PCR techniques can be utilized to introduce single amino acid substitutions at selected sites into gp120 fragments containing the neutralizing epitope. Briefly, overlapping portions of the region containing the epitope are amplified using primers that incorporate the desired nucleotide changes. The resultant PCR products are annealed and amplified to generate the final product. The final product is then expressed to produce a mutagenized gp120 fragment. Expression of DNA encoding gp120 or a portion thereof is described hereinafter and exemplified in the Examples.

In a preferred embodiment described in Example 1, the gp120 fragments are expressed in mammalian cells that are capable of expression of gp120 fragments having the same glycosylation and disulfide bonds as native gp120. The presence of proper glycosylation and disulfide bonds provides fragments that are more likely to preserve the neutralizing epitopes than fragments that are expressed in *E. coli*, for example, which lack disulfide bonds and glycosylation or are chemically synthesized which lack glycosylation and may lack disulfide bonds.

Those mutagenized gp120 fragments are then used in an immunoassay using gp120 as a control to determine the mutations that impair or eliminate binding of the neutralizing antibodies. Those critical amino acid residues form part of the neutralizing epitope that can only be altered in limited ways without eliminating the epitope. Each alteration that preserves the epitope can be determined. Such mutagenicity studies demonstrate the variations in the amino acid sequence

of the neutralizing epitope that provide equivalent or diminished binding by neutralizing antibodies or eliminate antibody binding. Although the amino acid sequence of gp120 used in the vaccine preferably is
5 identical to that of a selected HIV isolate for the given geographic region, alterations in the amino acid sequence of neutralizing epitope that are suitable for use in a vaccine can be determined by such studies.

Once a neutralizing epitope is localized to a
10 region of ten to twenty amino acids of gp120, the amino acid sequence of corresponding neutralizing epitopes of other HIV isolates can be determined by identifying the corresponding portion of the gp120 amino acid sequence of the isolate.

15 Once the neutralizing epitopes for a given region of gp120 are determined, the amino acid sequence of HIV isolates for the geographic region are determined. The complete amino acid sequence for numerous isolates has been determined and is available from numerous journal
20 articles and in databases. In such cases, determination of the amino acid sequence of HIV isolates for the geographic region involves looking up the sequence in an appropriate database or journal article. However, for some isolates, the amino acid
25 sequence information does not include the sequence of the V2 or C4 domains.

When the amino acid sequence of a region of interest for a given isolate is not known, the amino acid sequence can be determined by well known methods.
30 Methods for determining the amino acid sequence of a protein or peptide of interest are well known and are described in numerous references including Maniatis et al., Molecular Cloning--A Laboratory Manual, Cold Spring Harbor Laboratory (1984). In addition,
35 automated instruments which sequence proteins are commercially available.

Alternatively, the nucleotide sequence of DNA encoding gp120 or a relevant portion of gp120 can be determined and the amino acid sequence of gp120 can be deduced. Methods for amplifying gp120-encoding DNA from HIV isolates to provide sufficient DNA for sequencing are well known. In particular, Ou et al, *Science* 256:1165-1171 (1992); Zhang et al. *AIDS* 5:675-681 (1991); and Wolinsky *Science* 255:1134-1137 (1992) describe methods for amplifying gp120 DNA. Sequencing of the amplified DNA is well known and is described in Maniatis et al., *Molecular Cloning--A Laboratory Manual*, Cold Spring Harbor Laboratory (1984), and Horvath et al., *An Automated DNA Synthesizer Employing Deoxynucleoside 3'-Phosphoramidites*, *Methods in Enzymology* 154: 313-326, (1987), for example. In addition, automated instruments that sequence DNA are commercially available.

In a preferred embodiment, the isolate is a patient isolate which has not been passaged in culture. It is known that following passage in T-cells, HIV isolates mutate and isolates best suited for growth under cell culture conditions are selected. For example, cell culture strains of HIV develop the ability to form syncytia. Therefore, preferably the amino acid sequence of gp120 is determined from a patient isolate prior to growth in culture. Generally, DNA from the isolate is amplified to provide sufficient DNA for sequencing. The deduced amino acid sequence is used as the amino acid sequence of the isolate, as described hereinbefore.

To determine the percentage each isolate constitutes of total HIV that infects individuals in the geographic region, standard epidemiological methods are used. In particular, sufficient isolates are

sequenced to ensure confidence that the percentage of each isolate in the geographic region has been determined. For example, Ichimura et al, *AIDS Res. Hum. Retroviruses* 10:263-269 (1994) describe an
5 epidemiological study in Thailand that determined that there are two strains of HIV present in the region. HIV strains have only recently been present in Thailand and Thailand, therefore has the most homogenous population of HIV isolates known to date. The study
10 sequenced 23 isolates from various parts of the country and determined that only two different amino acid sequences were present in the isolates.

In contrast, HIV has been infecting individuals in Africa for the longest period of any geographic region.
15 In Africa, each of the most common isolates probably constitutes about 5% of the population. In such cases, more isolates would need to be sequenced to determine the percentage each isolate constitutes of the population. Population studies for determining the
20 percentage of various strains of HIV, or other viruses, present in a geographic region are well known and are described in, for example, Ou et al, *Lancet* 341:1171-1174 (1993); Ou et al, *AIDS Res. Hum. Retroviruses* 8:1471-1472 (1992); and McCutchan et al., *AIDS Res. Hum. Retroviruses* 8:1887-1895 (1992).
25

In the United States and western Europe, probably about two to four different neutralizing epitopes in each of the V2, V3, and C4 domains constitute 50 to 70% of the neutralizing epitopes for each domain in the
30 geographic region, as described more fully hereinafter.

Selection method

Once the amino acid sequence of neutralizing epitopes for strains in a region are determined, gp120
35 from an HIV strain having gp120 that has an amino acid sequence for a neutralizing epitope in the V2 or C4

domain which sequence is one of the most common in the geographic region is selected. One of the most common neutralizing epitope amino acid sequences means that the strain has an amino acid sequence for at least one neutralizing epitope that is occurs among the most frequently for HIV isolates in the geographic region and thus is present as a significant percentage of the population. For example, if there are three sequences for a neutralizing epitope that constitute 20, 30, and 40 percent of the sequences for that epitope in the region and the remainder of the population is comprised by 2 to 4 other sequences, the three sequences are the most common. Therefore, in African countries, if each of several amino acid sequences constitute about 5% of the sequences for a neutralizing epitope and the remainder of the sequences each constitute less than 1% of the population, the isolates that constitute 5% of the population are the most common.

Preferably, isolates having the most common amino acid sequences for a neutralizing epitope are chosen. By the most common is meant that the sequences occur most frequently in the geographic region. For example, in the United States, the MN isolate has a C4 neutralizing epitope that comprises at least about 45% of the population. The GNE₈ isolate has a C4 neutralizing epitope that comprises at least about 45% of the population. Thus either isolate has the most common C4 neutralizing epitope in the region. When gp120 from each isolate is combined in a vaccine, greater than about 90% of the C4 neutralizing epitope sequences are present in the vaccine. In addition, the amino acid sequences for the V3 neutralizing epitope in the MN and GNE₈ isolates are substantially similar and comprise about 60% of the population. Therefore, those strains have the two most common neutralizing epitopes for the V3 domain. In the V2 region, the MN isolate

amino acid sequences comprises about 10% of the population, and the GNE₃ isolate amino acid sequences comprises about 60% of the population. Therefore, the GNE₃ strain has the most common neutralizing epitope for the region and the two strains together comprise the two most common neutralizing epitopes for the region. A multivalent gp120 subunit vaccine containing the two isolates contains amino acid sequences for epitopes that constitute about 70% of the V2 domain, about 60% of the V3 domain, and about 90% of the C4 domain for the United States.

In a preferred embodiment of the method, one or more HIV isolates having an amino acid sequence for a neutralizing epitope in the V2 and/or C4 domains that constitute at least about 50% of the population for a selected geographic region are selected. In a more preferred embodiment, isolates having the most common neutralizing epitopes in the V3 domain are also included in the vaccine.

As is clear, once the most common amino acid sequences for the neutralizing epitopes in the V2, V3, and C4 domains are known, an isolate having a common epitope for each region is preferably selected. That is, when only two or three isolates are used for the vaccine, it is preferable to select the isolate for common epitopes in each region, rather than selecting an isolate by analysis of a single region.

In a more preferred embodiment, gp120 from isolates having epitopes that constitute at least 50% of the population for the geographic region for V2, V3, and C4 domains are present in the vaccine. More preferably, the isolates have epitopes that constitute at least 60% of the population for the geographic region for the three domains. Most preferably, 70% or more are included.

In another preferred embodiment, the entire amino acid sequence of the V2 and C4 domains is determined in the selection process. In addition to selecting common sequences for the neutralizing epitopes, isolates
5 having unusual polymorphisms elsewhere in the region are preferably not used for the vaccine isolates.

Vaccine preparation

gp120 from the selected HIV isolate(s) is used to
10 make a subunit vaccine, preferably a multivalent subunit vaccine. Preparation of gp120 for use in a vaccine is well known and is described hereinafter. With the exception of the use of the selected HIV isolate, the gp120 subunit vaccine prepared in the
15 method does not differ from gp120 subunit vaccines of the prior art.

As with prior art gp120 subunit vaccines, gp120 at the desired degree of purity and at a sufficient concentration to induce antibody formation is mixed
20 with a physiologically acceptable carrier. A physiologically acceptable carrier is nontoxic to a recipient at the dosage and concentration employed in the vaccine. Generally, the vaccine is formulated for injection, usually intramuscular or subcutaneous
25 injection. Suitable carriers for injection include sterile water, but preferably are physiologic salt solutions, such as normal saline or buffered salt solutions such as phosphate buffered saline or ringer's lactate. The vaccine generally contains an adjuvant.
30 Useful adjuvants include QS21 which stimulates cytotoxic T-cells and alum (aluminum hydroxide adjuvant). Formulations with different adjuvants which enhance cellular or local immunity can also be used.

Addition excipients that can be present in the
35 vaccine include low molecular weight polypeptides (less than about 10 residues), proteins, amino acids,

carbohydrates including glucose or dextrans, chelating agents such as EDTA, and other excipients.

The vaccine can also contain other HIV proteins. In particular, gp41 or the extracellular portion of gp41 can be present in the vaccine. Since gp41 has a conserved amino acid sequence, the gp41 present in the vaccine can be from any HIV isolate. gp160 from an isolate used in the vaccine can replace gp120 in the vaccine or be used together with gp120 from the isolate. Alternatively, gp160 from an isolate having a different neutralizing epitope than those in the vaccine isolates can additionally be present in the vaccine.

Vaccine formulations generally include a total of about 300 to 600 μ g of gp120, conveniently in about 1.0 ml of carrier. The amount of gp120 for any isolate present in the vaccine will vary depending on the immunogenicity of the gp120. For example, gp120 from the Thai strains of HIV are much less immunogenic than gp120 from the MN strain. If the two strains were to be used in combination, empirical titration of the amount of each virus would be performed to determine the percent of the gp120 of each strain in the vaccine. For isolates having similar immunogenicity, approximately equal amounts of each isolate's gp120 would be present in the vaccine. For example, in a preferred embodiment, the vaccine includes gp120 from the MN, GNE₈, and GNE₁₆ strains at concentrations of about 300 μ g per strain in about 1.0 ml of carrier. Methods of determining the relative amount of an immunogenic protein in multivalent vaccines are well known and have been used, for example, to determine relative proportions of various isolates in multivalent polio vaccines.

The vaccines of this invention are administered in the same manner as prior art HIV gp120 subunit

vaccines. In particular, the vaccines are generally administered at 0, 1, and at 6, 8 or 12 months, depending on the protocol. Following the immunization procedure, annual or bi-annual boosts can be administered. However, during the immunization process and thereafter, neutralizing antibody levels can be assayed and the protocol adjusted accordingly.

The vaccine is administered to uninfected individuals. In addition, the vaccine can be administered to seropositive individuals to augment immune response to the virus, as with prior art HIV vaccines. It is also contemplated that DNA encoding the strains of gp120 for the vaccine can be administered in a suitable vehicle for expression in the host. In this way, gp120 can be produced in the infected host, eliminating the need for repeated immunizations. Preparation of gp120 expression vehicles is described hereinafter.

20 Production of gp120

gp120 in the vaccine can be produced by any suitable means, as with prior art HIV gp120 subunit vaccines. Recombinantly-produced or chemically synthesized gp120 is preferable to gp120 isolated directly from HIV for safety reasons. Methods for recombinant production of gp120 are described below.

DNA Encoding GNE₈ and GNE₁₆ gp120 and the resultant proteins

30 The present invention also provides novel DNA sequences encoding gp120 from the GNE₈ and GNE₁₆ isolates which can be used to express gp120 and the resultant gp120 proteins. A nucleotide sequence of less than about 5 kilobases (Kb), preferably less than about 3 Kb
35 having the nucleotide sequence illustrated in Tables 1 and 2, respectively, encodes gp120 from the GNE₈ and

GNE₁₆ isolates. The sequences of the genes and the encoded proteins are shown below in Tables 1-3. In particular, Table 1 illustrates the nucleotide sequence (SEQ. ID. NO. 27) and the predicted amino acid sequence (SEQ. ID. NO. 28) of the GNE₁₆ isolate of HIV. The upper sequence is the coding strand. The table also illustrates the location of each of the restriction sites.

hgiCI
 banI
 bspl286
 bmyI
 styI
 scfI
 bsgI
 pstI
 scfI

1 ATGATAGTGA AGGGGATCAG GAAGAATTGT CAGCATTGT GGAGATGGG CACCATGCTC CTGGGATGT TGATGATCTG TAGTGCTGCA GAAAAATGT
 TACTATCACT TCCCCTAGTC CTTCTTARCA GTCGTGRACA CCTCTACCCC GTGCTACGAG GAACCTTACA ACTACTAGAC ATCAGGACCT CTTTTTACA
 1 M I V K G I R K N C Q H L W R W G T M L L G M L M I C S A A E K L W

kpnI
 hgiCI
 banI
 asp718
 acc651
 ndeI

101 GGGTCACAGT CTATTATGGG GTACCTGTGT GGAAGAGACC AACCACTACT CTAATTGTG CATCAGATGC TAAAGCATAT GATACAGAGG TACATAATGT
 CCCAGTGTCA GATAATACCC CATGGACACA CTTTCTCTG TTGGTGTGA GATAAACAC GTAGTCTACG ATTTCGTATA CTATGCTCC ATGTATTACA
 35 V T V Y Y G V P V W K E A T T T L F C A S D A K A Y D T E V H N V

nspl
 nsphi
 nsphi
 apoI
 aflIII

201 TTGGGCCACA CATGCTGTG TACCCACAGA CCCCAACCCA CAAGAATAG GATTGGAAGA TGTAAACAGAA AATTTTAACA TGTGGAAGA TAACATGGTA
 AACCGGTGT GTACGGACAC ATGGGTGTCT GGGGTGGGT GTTCTTTATC CTAACCTTTT ACATGTCTTT TTAACATGT ACACCTTTT ATGTACCAT
 68 W A T H A C V P T D P N P Q E I G L E N V T E N F N M W K N N M V

ppulOI
 nsII/avaIII
 hindIII
 draIII
 ahaIII/draI

301 GAACAGATGC ATGAGGATAT AATCAGTTTA TGGGATCAAA GCTTAARACC ATGTGTAAAA TTAACCCAC CATGTGTTAC TTAAATTCG ACTGATTGA
 CTTGCTAGC TACTCCTATA TTAGTCAAT ACCCTAGTT CGAATTGGG TACACATTTT AATTGGGGT ATACACAATG ARAATTACG TGACTAACT
 101 E Q M H E D I I S L W D Q S L K P C V K L T P L C V T L N C T D L K

pvuII
 speI
 nsbII
 scfI

401 AARATGCTAC TAATPACCACT ACPAGCAGCT GGGGAARAGT GGAGAGAGGA GAATAAAAA ACTGCTCTTT CAATGTCACC ACAAGTATAA GAGATAAGAT
 TTTTACGATG ATTATGGTGA TCATCGTGA CCCCTTTCTA CCTCTCTCT CTTTATTTT TGACAGAGAA GTTACAGTGG TGTTCATATT CTTATTTCTA
 135 N A T N T T S S S W G K M E R G E I K N C S F N V T T S I R D K M

nspl
 nsphi
 nsphi
 apoI
 aflIII

501 GAAGAATGAA TATGCACTTT TTTATAAAT TGATGTAGTA CCAATAGATA ATGATATATAC TAGCTATAGG TTGATRAAGT GTACACCTC AGTCATTACA
 CTTCTACTT ATACGTGAAA AATATTGTA ACTACATCAT GGTATCTAT TACTATTATG ATCGATATCC AACTATTCAA CATTTGGAG TCAGTAACT
 168 K N E Y A L F Y K L D V V P I D N D N T S Y R L I S C N T S V I T

stuI
 ha I
 bsp1286
 bmyI
 601 CAGGCCCTGTC CAAGGCTGTC CTTTGAGCCA ATTCCCATAC ATTATTGTGC CCGGCTGGT TTTGGATTG TAAAGTGTAG AGATAAAAG TTCACCGGAA
 GTCCGGACAG GTTTCACAG GAACTCGGT TAAGGTATG TAATAACACG GGGCGGACCA AAACGCTAAG ATTACATAC TCTATTTTC AAGTTGCCCT
 201 Q A C P K V S F E P I P I H Y C A P A G F A I L K C R D K K F N G T
 bsp14071
 haeI
 701 CAGGACCATG TACAAATGTC AGCACAGTAC TCGTGTATG TTACATGTGT ACCITTAATCC GGTCAATGTA GTTGAATGA CGACAAATTA CCGTCAATC GTCCTCTCT
 GTCTGTGATC ATGTTTACAG TCGTGTATG TCGTGTATG TCGTGTATG TCGTGTATG TCGTGTATG TCGTGTATG TCGTGTATG TCGTGTATG TCGTGTATG
 235 G P C T N V S T V Q C T H G I R P V V S T Q L L L N G S L A E E E
 bsp14071
 pvuII
 bstXI/xhoII
 bglII
 801 AGTAGTAAT AGATCTGCA ATTCTCGCA CAATGCTAAG ACCATAATAG TACAGCTGAA CGAATCTGTA GAAATTAAT GTACAGAGCC CAACAACAT
 TCATCATTA TCTAGACGCT TAAGAGCCCT GTTACGATTT TGGTATATC ATGTGCACTT GTTAGACAT CTTAATTA CATGTTCTGG GTTGTGTTA
 268 V V I R S A N F S D N A K T I I V Q L N E S V E I N C T R P N N N
 bst11071
 accI
 901 ACAAGAAGAA GTATACATAT AGGACCAGG AGAGCATTTT ATGCAACAGG AGAATAATA GGACACATAA GACACACACA TTGTAACCTT AGTACACAA
 TGTCTTCTT CATATGATA TCTGTTGCCC TCTGTTGAAA TACGTTGTC TCTTATAT CTTCTGTTT CTTCTGTTT CTTCTGTTT CTTCTGTTT CTTCTGTTT
 301 T R R S I H I G P G R A F Y A T G E I I G D I R Q A H C N L S S T K
 ahaIII/draI
 1001 AATGGAATA TACTTTAAA CAGATAGTTA CAAATTAAG AGACATTTT AATAAACA TAGTCTTAA TCACTCTCA GGAGGGACCC CAGAAATTTG
 TTACTTATT ATGAATTTT GTCTATCAAT GTTTTAATC TCTTGTAAA TTATTTTGT ATCAGAAAT AGTGAGAGT CCTCCCTGG GTCTTTACA
 335 W N N T L K Q I V T K L R E H F N K T I V F N H S S G G D P E I V
 apoI
 1101 AATGCACAGT TTTAATTGT GAGGGGAAT TTCTACTGT AATACAAC CACTGTTTAA TAGTACTTGG AATTATCTT ATACTTGGAA TAATCTGAA
 TTACGTGTCA AATTAACAC CTCCCTTAA AAGATGACA TTATGTTGT GTGACAAAT ATCATGACCC TTAATATGAA TATGACCTT ATTATGACTT
 368 M H S F N C G G E F F Y C N T T P L F N S T W N Y T Y T W N N T E
 scaI
 1201 GGTCAATG AACTGGAAG AATATCACA CTCCAATGCA GAATAAACA AATTATAC ATGTGGCAGG AAGTAGGAA AGCAATGTAT GCGCTCCCA
 CCGAGTTAC TGTACCTTC TTTATAGTGT GAGTTACGT CTATTTTGT TTAATATTG TACACCGTCC TCACTCTT TCGTTACATA CGGGAGGCT
 401 G S N D T G R N I T L Q C R I K Q I I N M W Q E V G K A M Y A P P I
 nspl
 nspl
 afIII
 eco571

1301 TAAGAGGACA AATTAGATGC TCATCAATA sspl
 435 ATTCTCCTGT TTAATCTAGG AGTAGTTTAT S S N I T G L L L T R D G G N N S E T E I F R P G G G
 1401 AGATATGAGG GACAATTGGA GAAGTGATT ATATAAATAT AAAGTAGTAA AATTTGAACC ATTAGGAGTA GCACCCACCA AGCAAAAGAG AAGAGTGATG
 468 TCTATACTCC CTGTTAACCT CTTCACTTAA TATATTATA TTTCACTAAT TTTACTTGG TAATCCTCAT CGTGGGTGGT TCCGTTTCTC TTCTACATAC
 1501 CAGAGAGAAA AAGAGCAGT GGAATAGGA GCTGTGTTCC TTGGTTCTT GGGAGCAGCA GGAAGCACTA TGGGCGCAGC GTCAGTGAGC CTGACGGTAC
 501 GTCTCTCTTT TTCTCGTCA CCCTTATCCT CGACACAAGG AACCCAGAA CCCTCGTCTG G A A G S T M G A A S V T L T V Q
 1601 AGCCAGACT ATTATTGTCT GGTATAGTGC AACAGCAGAA CAATTGCTG AGGGTATTG AGGCCAACA GCATCTGTTG CAATCAGACG TCTGGGGCAT
 535 TCCGGTCTGA TAATAACAGA CCATATCAGG TTGTGCTCTT GTTAACGAC TCCCGATAAC TCCGGCTGTG CGTAGACAAAC GTTGAAGTGC AGACCCGTA
 1701 CRAGCAGCTC CAGCAGAG TCCTGGCTGT GGAGAGATAC CTAAGGATC AATAGGATTC GGGGATTGG GGTGCTCTG GAAACTCAT CTGCACTACT
 568 GTTGGTCTGA GTCCGTTCTC AGGACCGACA CCTCTCTATG GATTTCTATG TTGTCGAGGA CCCCTAACC CCAACGAGAC CTTTTCAGTA GACGTGTA
 1801 GCTGTGCTCT GGAATGCTAG TTGGAGTAAT AAATCTCTGG ATAAAGATTG GGATAACATG ACCTGGATGG AGTGGGAAG AGAATTGAC AATTACACA
 601 CGACACGGAA CCTTACGATC AACCTCATTA TTAGAGACC TATTCTAAC CCTATTGTAC TGGACCTTAC TCACCTTTC TCTTTAAGT TTAATGTGT
 1901 GCTTAATATA CAGCTTAATT GAAGAATCGC AGAACCAACA AGAAAAAAT GAACAGAAT TATTGGAAT AGATAAATGG CCAAGTTTCT GGAATTGTT
 635 CGAATTATAT GTCGAATTA CTTCTTAGCG TCTTGGTTGT TCTTTTTTA CTGTCTTCTTA ATAACCTTAA TCTATTACC CGTTCAACA CTTAACCNA
 2001 TGACATAACA AAATGGCTGT GGTATATAA AATATTCATA ATGATAGTAG GAGGCTTGGT AGGTTTAAAG ATAGTTTTTA CTGTACTTTC TATAGTGAAT
 668 ACTGTATTGT TTTACCGACA CCATATATT TTAAGAAT TACTATCATC CTCCGACCA TCCAAATCT TATCAAAAT GACATGAAG ATATCACTTA

eco571 bstyl/xhoII gsuI/bpmI
 bgIII econI
 styI earI/ksp6321
 muni
 hael
 alwNI
 alwNI
 alwNI
 gsuI/bpmI
 bstyl
 bsmI
 hindIII
 sspi
 scfI

30

Table 2 illustrates the nucleotide sequence and the predicted amino acid sequence of the GNE₁₆ isolate of HIV. The upper sequence is the coding strand. The table also illustrates the location of each of the restriction sites. The first four pages of the table are from one clone of the gene and the second three pages of the table are from another clone of the gene. The sequences of the clones differ by about 2%. (The nucleotide sequences are SEQ. ID. NOs. 28 and 29, respectively. The amino acid sequences are SEQ. ID. NOs. 30 and 31, respectively.) It is noted that each of the sequences includes a stop codon. A gene sequence that encodes full length gp120 can be made by repairing one of the sequences.

15

TABLE 2

1	ATGAGAGTGA AGGGATCAG GAGGAATTAT CAGCAGCTTGT GGAGATGGGG CACCATGCTC TGATGATCTG TAGTCTCTGA GGGAAATTGT	scfi
	TACTCTCACT TCCCCTAGIC CTCCTTAATA GTCGTGAACA CCTCTACCC GTGGTACGAG GRACCTATA ACTACTAGAC ATACGACGT CCGTTAACA	psfi
1	M R V K G I R R N Y Q H L W R W G T M L L G I L M I C S A A G K L W	bsgi
		scfi
		styI
		bmyI
		bsp1286
		hgiCI
		banI
		kpnI
		hgiCI
		banI
		asp718
		acc65I
101	GGGTCACAGT CTATTATGGG GTACCTGTGT GGAAGAAAC AACCCACCACT CTATTTTGTG CATCAGATGC TAAAGCATAT GATACAGAGA TACATATGT	ndel
	CCAGTGTCA GATAATACCC CATGGACACA CCTTCTTTG TTGGTGGGA GATAAAGAC GTAGCTACG ATTCGTATA CTATCTCT ATGTATTACA	
35	V T V Y Y G V P V W K E T T T T L F C A S D A K A Y D T E I H N V	
		nspl
		nsphi
		afIII
		apoi
201	TTGGGCCACA CATGCTGTG TACCCACAGA CCCCAACCCA CAAGAAGTAG TATTGGAAA TGTGACAGAA AATTTTACA TGTGGAATAA TAACATGTG	
	AACCGGTGT GTACGACAC ATGGGTGTCT GGGTGGGT GTTCTTCATC ATAACCTTTT ACACGTCTT TTAATAATT ACACCTTTT ATTGTACCAC	
68	W A T H A C V P T D P N P Q E V V L E N V T E N F N M W K N N M V	
		nspl
		nsphi
		afIII
		ahaiII/drai
		ahaiII/drai
301	GAACAGATGC ATGAGATAT AATCAGTTA TGGGATCAA GTTAAAGCC ATGTGTAATA TTAACCCAC TCTGTGTAC TTAAATTGC ACTGATGCGG	
	CTTGCTACG TACTCTATA TTAGTCAAT ACCTAGTTT CAAATTGCG TACACATTTT AATTGGGGT AGACACAAAG AAATTTAAG TGACTACGCC	
101	E Q M H E D I I S L W D Q S L K P C V K L T P L C V T L N C T D A G	
		gsul/bpmI
401	GGAACTACTA TAATACCAAT AGTAGTAGCA GGGAAAAGCT GGGAAGAAGG GAAATAAAA ACTGCTCTTT CAATATCACC ACAAGCGTGA GAGATAAGAT	
	CCATTAGATG ATTATGGTTA TCATCATCGT CCTTTTCTG CCTTTTCTT CCTTTTCTT TGACGAGAAA GTTATAGTGG TGTTCGCACT CTCTATTCTA	
135	N T T N T N S S S R E K L E K G E I K N C S F N I T T S V R D K M	
		scal
		scfi
501	GCAGAAAGAA ACTGCACCTT TTAATAAAGT TGATATAGTA CCAATAGATG ATGATAGTAG GAATAGTACT AGGAATAGTA CTAACATAG GTTGATAAGT	
	CGTCTTTCTT TGACGTGAAA AATTATTGA ACTATATCAT GGTATCTAC TACTACTATC CTTATCTATG TCCCTATCAT GATTGATATC CAACATATTCA	
168	Q K E T A L F N K L D I V P I D D D R N S T R N S T N Y R L I S	
		^421,reverse
		^43r2,reverse

stul
 haeI
 601 TGTAAACACCT CAGTCATTAC ACAGGCCCTGT CCRAAGGTAT CATTTCAGCC AATTCACATA CATTCTCTGA CCCGGCTGG TTTTGGCTT CTAAAGTGA
 ACATTGTGGA GTCAGTAATG TGTCCGGACA GGTTCACATA GTAACTCGG TTAAGGGTAT GTAAAGACAT GGGCCGACC AAAACGGAA GATTCACAT
 201 C N T S V I T Q A C P K V S F E P I P I H F C T P A G F A L L K C N
 bsp1407I haeI
 701 ATAAATAGAC GTTCAATGGA TCAGGACCAT GCAAAATGT CAGCACAGTA CAATGTACAC ATGAATAG GCGAGTAGTA TCAATCAAC TCGTGTAA
 TATTATCTG CAAGTTACCT AGTCCTGGTA CGTTTTTACA GTCGTGTCAT GTTACATGTG TACCTTAATC CGGTCATCAT AGTTAGTTG ACACAATTT
 235 N K T F N G S G P C K N V S T V Q C T H G I R P V V S T Q L L L N
 bstYI/xhoII
 801 TGGCAGTCTA GCGAAGGAG AGGTAGTAAT TAGATCTGAA AATTTCACGA ACAATGCTAA AACCATAATA GTACAGCTGA CAGACCCAGT AAAAATAAT
 ACCGTCAGAT CGCTTCTC TCATCATTA ATCTAGACTT TTAAGTGCT TGAAGTGTG TGTACGATTT TGGGTATAT CATGTCGACT GTCTTGTCA TTTTAAATTA
 268 G S L A E G E V I R S E N F T N A K T I I V Q L T E P V K I N
 ^fl, forward
 bst1107I
 901 TGTACAGAC CCAACAACAA TACAAGAAA AGTATACCTA TAGGACCAGG GAGAGCATTT TATGCAACAG GAGACATAAT AGGAATATA AGACAGCAC
 ACATGTTCTG GGTGTTGTT ATGTTCTTTT TCATATGGAT ATCCTGGTCC CTCTCGTAAA ATACGTTGTC CTCTGTATTA TCCTTTTAT TCTGTTCTG
 301 C T R P N N N T R K S I P I G P G R A F Y A T G D I I G N I R Q A H
 ^875, reverse
 bst1107I
 1001 ATTGTRACCT TAGTGAACA GACTGGAATA ACACTTTAGG ACAGATAGTT GAAAAATTA GAGACAATTT TGGGAATRAA ACAATAATCT TTAATCACTC
 TAACATTGGA ATCATCTGT CTGACCTTAT TGTGAATCC TGTCTATCAA CTITTTAAT CTCTGTAA ACCCTTATTT TGTATTAGA AATTAGTGAG
 335 C N L S R T D W N N T L G Q I V E K L R E Q F G N K T I I F N H S
 ppuMI
 eco0109I/draII
 1101 CTCAGGAGG GACCCAGRA TTGTAATGCA CAGTTTAAAT TGTAGAGGG AATTTTCTA CTGTAATACA ACACAATTTG TTGACAGTAC TTGGGATAAT
 GAGTCTCCC CTGGGTCTTT AACATTAGT GTCAAAATTA ACATCTCCC TTAAGAGAT GACATTATGT TGTGTTAACA AACTGTCATG AACCTTATTA
 368 S G G D P E I V M H S F N C R G E F F Y C N T T Q L F D S T W D N
 earI/ksp632I
 eco57I
 1201 ACTAAAGTGT CAAATGGCAC TAGCACTGAA GAGAATAGCA CAATCAGCT CCCATGCAGA ATAAAGCAA TTGTAACAT GTGCGAGAA GTAGGAAAG
 TGATTTCACA GTTACCGTG ATCGTGACT CTCTTATCGT GTTAGTGGA GGTAGCTCT TATTGCTTT AACATTGTA CACGTCCTT CATCCTTTT
 401 T K V S N G T S T E E N S T I T L P C R I K Q I V N M W Q E V G K A

eco81I
 bsu36I/
 mstII/
 sauI

muni
 scaI

nspI
 nsphi
 afIII

1301 CAATGATGCG CCTCCCATC AGAGGACAAA TTAGATGCTC ATCAATATT ACAGGTTGC TATTACAAG AGATGGAGGT AGTACAACA GCATGAATGA ^{bsal}
 GTTACATACG GGGAGGGTAG TCTCCTGTTT AATCTACAAG TAGTTTATAA TGTOCCAACG AATATTGTTT TCTACCTCCA TCATTGTTGT CGTACTTACT
 435 M Y A P P I R G Q I R C S S N I T G L L L T R D G G S N S M N E
 ^2,16.7f3,forward

1401 GACCTTCAGA CCTGGAGGAG GAGATATGAG GGACAAATGG AGAAGTGAAT TATACAATA TAAAGTAGTA AAAATTGAAC CATTAGGAGT AGCACCACCC ^{styI}
 CTGGAAGTCT GGACCTCCTC CTCATATCTC CCGTTTAACC TCTTCACTTA ATATGTTTAT ATTCATCAT TTTAACCTG GTAATCCTCA TCGTGGGTGG
 468 T F R P G G G D M R D N W R S E L Y K Y K V V K I E P L G V A P T
 ^c4rev4,reverse

1501 AAGGCAAGA GAACAGTGGT GCAGAGAGAA AAAAGAGCAG TGGGAATAGG AGCTGTGTTT CTGGGTTCT TAGGAGCAGC AGGAAGCACT ATGGGCGCAG ^{styI}
 TTCCGTTTCT CTCTCACCA CGTCTCTCTT TTTTCTCCTC ACCCTTATCC TCGACACAAG GAACCAAGA ATCTCGTGC TCCTTCGTGA TACCCGCGTC
 501 K A K R R V V Q R E K R A V G I G A V F L G F L G A A G S T M G A A
 haeI

1601 CGTCAATAAC GCTGACGGTA CAGGCCAGAC TATTATTGTC TGGTATAGTG CAACAGCAGA ACATTTGCT GAGGCTATT GAGGCGAAC AGCATCTGTT ^{alwNI}
 GCAGTTATTG CGACTGCCAT GTCCGGTCTG ATAATAACAG ACCATATCAC GTTGCTGCTT TGTAAACGA CTCGCGTTG TCCTAGACAA
 535 S I T L T V Q A R L L L S G I V Q Q Q N N L L R A I E A Q Q H L L
 ^43f5,forward

1701 GCAACTCATA GTCTGGGGCA TCAAGCAGCT CCAGGCAAGA GTCCTGGCTG TGGAAACATA CCTAAGGGAT CAACAGCTCC TGGGATTGG GGGTTGCTCT ^{eco81I}
 CGTTGAGTAT CAGACCCCGT AGTTGTCGA GGTCCGTTCT CAGGACCGAC ACCTTCTAT GGAATCCCTA GTTGCGAGG ACCCTAAAC CCCAAGGAGA ^{bsu36I/mstII/sauI}
 568 Q L I V W G I K Q L Q A R V L A V E R Y L R D Q Q L L G I W G C S

1801 GGAACAATCA TTTCACCCAC CTCAGTGCCT TGAATGCTA GTTGAGTAA TAAATCTCTA GATAAGATT GGAATACAT GACCTGGATG GAGTGGGAAA ^{xbaI}
 CCTTTGAGT AAACGTGGTG GAGTCACGGA ACCTTACGAT CAACCTCAT ATTAGAGAT CTATTTCTAA CCCTATTGTA CTGGACCTAC CTCACCTTT
 601 G K L I C T T S V P W N A S W S N K S L D K I W D N M T W M E W E R

1901 GAGAAATGA GAATTACAC AGCTTAATAT ACACCTTAAT TGAAGAATCG CAGAACCAAC AGAARAGAA TGAACAGAC TTATGGAAAT TGGATCAATG ^{hindIII}
 CTCTTAACT CTTAATGTT TCGAATTATA TGTGGAATTA ACTTCTTAGC GTCTTGGTTG TTCTTTCTT ACTTGTCTG AATAACCTTA ACCTAGTTAC
 635 E I E N Y T S L I Y T L I E E S Q N Q Q E K N E Q D L L E L D Q W

2001 GGCAAGTCTG TGGATTGGT TTAGCATTAAC AAATGGCTG TGGTATATAA AATATTCAT AATGATAGTT GGAGGCTGG TAGGTTTAAAG AATAGTTTCTT
 CCGTTCAGAC ACCTTAACCA AATCGTATTG TTTTACCGAC ACCATATATT TTATATAAGTA TTACTATCAA CTCCCGAACC ATCCAATTC TTATCAAAAA
 668 A S L W N W F S I T K W L W Y I K I F I M I V G G L V G L R I V F
 ^43f6, forward ^2000, reverse
 scfI
 2101 GCTGTACTTT CTATATGAA TAGAGTTAGG CAGGATTAAT CACCATATATC GTTTCAGACC CGCTCCCG CCGGAGGAG ACCCGACAGG CCCGAAGGAA
 CGACATGAAA GATATCACTT ATCTCAATCC GTCCCTATGA GTGGTAATAG CAAAGTCTGG GCGGAGGTC GGGCTCCTC TGGGCTGTCC GGGCTTCTT
 701 A V L S I V N R V R Q G Y S P L S F Q T R L P A P R R P D R P E G I
 xcmI
 2201 TCGAAGAGA AGGTGGAGAG CAAGGCAGAG ACAGATCCAT TCGCTTAGTG GATGATTCT TAGCACTTAT CTGGGAGGAC CTACGGAGCC TGTGCTCTT
 AGCTTCTTCT TCCACCTCTC GTTCCGCTC TGTCTAGTA AGCGAATCAC CTACCTAAGA ATCTGAATA GACCTGCTG GATGCTCGG ACACGGAGAA
 735 E E E G G E Q G R D R S I R L V D G F L A L I W D D L R S L C L F
 ^rl, reverse
 scfI
 2301 CAGCTACCAC CGCTTGAGAG ACTTACTCTT GATTGCAAG AGGATTGTGG AACTTGTGG AGCAGGGGG TGGGAAGCCC TCAATATTG GTGAATCTC
 GTCGATGGTG GCGAATCTC TGAATGAGAA CTAAGCTTGC TCCTAACACC TTGAAGACCC TCGTCCCCC ACCCTTCGG AGTTATAAC CACCTTAGAG
 768 S Y H R L R D L L L I A T R I V E L L G R R G W E A L K Y W N L
 scfI
 2401 CTACAGTATT GGATTGAGG ACTAAGAAAT AGTGTGTAA GCTTGCTTAA TGTACAGACC ATAGCAGTAG CTGAGGGGAC AGATAGGTT TTAGAAGTAT
 GATGTCATAA CCTAAGTCTT TGATTCTTA TCACGACAAAT CGAACGAAT ACAGTGTGG TATCGTATC GACTCCCTG TCTATCCAA AATCTCATA
 801 L Q Y W I Q E L K N S A V S L L N V T A I A V A E G T D R V L E V L
 2501 TACAAGAGC TTATAGAGCT ATTCTCCACA TACCTACAAG AATAGACAG GCGTTGGAA GGGCTTCTCT ATAA
 ATGTTCTCG AATATCTGA TAAGAGGTGT ATGATGTTC TTATCTGTC CCGAACCTTT CCCGAAGCA TATT
 835 Q R A Y R A I L H I P T R I R Q G L E R A L L O

ear1/ksp632I
 1 ATGAGAGTGA AGAGGATCAG GAGGAATTAT CAGCACTTCT GGAATGGG CACCATGCTC CTGGGATGT TGATGATCTG TAGTGCTGCA GGAATATTGT
 TACTCTCACT TCTCCTAGTC CTCCTTAATA GTCGTGAACA CCTTTACCC GGTGTACGAG GAACCTTACA ACTACTAGAC ATCAGCAGCT CCTTTTAA
 1 M R V K R I R R N Y Q H L W K W G T M L L G M L M I C S A A G K L W

hgiCI
 bani
 bsp1286
 bmyI
 kpn I
 hgiCI
 bani
 asp718
 acc65I
 101 GGGTCACAGT CTATTATGGG GTACCTGTGT GGAAGAAGAAC AACCACTCT CTATTGTGT CATCATGTC TAACGATAT GATACAGAGA TACATAATGT
 CCCAGTGTCA GATAATACCC CATGGACACA CCTTCTTTG TTGGTGTGA GATAAACAC GTAGCTACG ATTTCGTATA CTATGCTCT ATGTATTACA
 35 V T V Y Y G V P V W K E T T T T L F C A S D A K A Y D T E I H N V

nspl
 nsphi
 201 TTGGGCCACA CATGCCCTGT TACCCACAGA CCCCAACCCA CAAGAAGTAG TATTGGRAA TGTGACAGAA AATTTTAAACA TGTGGAATAA TAACATGCTG
 AACCCGGTGT GTACGGACAC ATGGGTGTCT GGGGTGGGT GTTCTTCATC ATACCTTTT ACACGTCTT TTAATATTGT ACACCTTTT ATTGTACCA
 68 W A T H A C V P T D P N P Q E V V L E N V T E N F N M W K N N M V

ppul01
 nsII/avaIII
 301 GAACAGATGC ATGAGGATAT AATCAGTTTA TGGGATCAA GTCTAAGCC ATGTGTAAA TTAACCCAC TCTGTGTAC TTTAATATGC ACTGATGCGG
 CTTCTCTACG TACTCCTATA TTACTCAAT ACCTAGTTT CAGATTTCG TACACATTT AATTGGGGT AGACACATG AATTTAAG TGACTACGCC
 101 E Q M H E D I I S L W D Q S L K P C V K L T P L C V T L N C T D A G

gsuI/bpmI
 401 GGAATACAT TAATACCAAT AGTACTAGCG GGGAAGAGCT GGAGAAAGGA GAAATAAAA ACTGCTCTTT CAATATCACC ARAAGCATGA GAGATAAGAT
 CCTATGATG ATTATGGTTA TCATCATCGC CCTTTTCGA CCTCTTCTT CTTTATTTT TGACGAGAAA GTTATAGTGG TGTTCTGACT CTCATTTCA
 135 N T T N T N S S S G E K L E K G E I K N C S F N I T T S M R D K M

scaI
 scaI
 501 GCAGAGAGAA ACTGCACCTT TTAATAAACT TGATATAGTA CCAATAGATG ATGATGATAG GAATAGTACT AGGAATAGTA CTAATATAG GTTGATAAGT
 CGTCTCTCT TGACGTGAAA AATTATTGA ACTATATCAT GGTATCTAC TACTACTATC CTATCATGA TCCTTATCAT GATTGATATG CAACATATCA
 168 Q R E T A L F N K L D I V P I D D D D R N S T R N S T N Y R L I S

stul
 haeI
 601 TGTAACACCT CAGTCATTAG ACAGGCTGT CCAAGGTAT CATTGAGCC AATTCCTATA CATTCTGTA CCCCGGCTGG TTTGGGCTT CTAAGTGTGA
 ACATTGTGA GTCAGTAATG TGTCGGACA GGTTCCTATA GTAACCTCG TTAAGGTAT GTAAAGACAT GGGCCGACC AAAACGGGA GATTTCACAT
 201 C N T S V I T Q A C P K V S F E P I P I H F C T P A G F A L L K C N

esp31
 701 ATATGAGAC GTTCAATGGA TCAGGACCAT GCAAAATGT CAGCACAGTA CTATGTACAC ATGGAATAG GCCAGTAGTA TCAACTCAAC TCGTGTAAA
 235 TATTACTCTG CAGTTACCT AGTCTGGTA CGTTTACGTA GCGTGTGCAT GATACATGTG TACCTTAATC CCGTCATCAT AGTTGAGTTG ACGACAATTT
 N E T F N G S G P C K N V S T V L C T H G I R P V V S T Q L L L N
 scaI bsp1407I haeI
 bstYI/xhoII
 earI/ksp32I
 801 TGGCAGTCTA GCAGGAGAAG AGGTAGTAAT TAGATCTGAA AATTTCACGA ACAATGCTAA AACCATATA GTACAGCTCA AAGAACCGT AAAAATTAAT
 ACCGTCAGAT CGTCTCTTC TCCATCATT ATCTAGACTT TTAAGTGTCT TGAAGTGTCT TGTAGGATTT CATGTCGAGT TTCTTGGTGA TTTTAAATTA
 268 G S L A G E E V I R S E N F T N N A K T I I V Q L K E P V K I N
 bst1107I
 accI scfI
 901 TGTACAGAC CCAACAACAA TACAAGAAA AGTATACCTA TAGGACCAGG GAGAGCATTT TATGCAACAG CGGACATAAT AGGAATATA AGACAAGCAC
 ACATGTTCTG GGTGTTGTT ATGTTCTTTT TCATATGAT ATCTGGTCC CTCTCGTAA ATACGTTGTC CGCTGTATTA TCCTTTATAT TCTGTTGCTG
 301 C T R P N N N T R K S I P I G P G R A F Y A T G D I I G N I R Q A H
 ec 81I
 bau361/
 metII/
 saul
 1001 ATTGTAACCT TAGTAGACA GACTGGATA ACACITTAAG ACAGATAGCT GAAATTAAT GAAACAATTT TGGGAATAAA ACAATAATCT TTAATCACTC
 TAACATTTGA ATCATCTTGT CTGACCTTAT TGTGAATTC TGTCTATCGA CTTTAAAT CTTTGTAA ACCCTTATTT TGTATTAGA AATTAGTAG
 335 C N L S R T D W N N T L R Q I A E K L R K Q F G N K T I I F N H S
 ppuNI
 eco0109I/draII
 1101 CTCAGGAGG GACCCAGAAA TTGTAATGCA CAGTTTAAAT TGTAGAGGG AATTTTCTA CTGTGATACA ACACAATTTGT TTAACAGTAC TTGGAATGCA
 GAGTCTCCC CTGGTCTTT AACATTACGT GTCAAAATTA ACATCTCCC TTAAGAAGAT GACACTATGT TGTGTAACA AATTGTCATG AACCTTACGT
 368 S G G D P E I V M H S F N C R G E F F Y C D T T Q L F N S T W N A
 nspl
 nsphI
 aflIII
 1201 AATAACACTG AAGGAATAG CACTAAGAG AATAGCACAA TCACACTCCC ATGCAGATA AAACAATTTG TAACATGTTG GCAGGAAGTA GGAAGAAGCA
 TTATTGTGAC TTTCCCTATC GTGATTTCTC TTATCGTGT AGTGTGAGG TACGCTTAT TTTGTTTAAAC ATTTGTACAC CGTCTTCAT CTTTTCGTT
 401 N N T E R N S T K E N S T I T L P C R I K Q I V N M W Q E V G K A M
 mamI
 bsaBI
 sspI
 1301 TGTATGCCC TCCCATCAGA GCACAATTA GATGTTTATC AATATTACA GGTGTTGCTAT TAACAAGAGA TGGAGGTAGT AGCAACAGCA TGAATGAGAC
 ACATACGGG AGGCTAGTCT CCTGTTTAAAT CTACAGTAG TTTATATATG CCAACAGATA ATTGTTCTCT ACCTCCATCA TCGTTCTCGT ACTTACTCTG
 435 Y A P P I R G Q I R C S S N I T G L L L T R D G G S S N S M N E T
 bsaI

eco57I econI gsuI/bpmI muniI styI styI
 1401 CTTACAGACCT GGAGGAGGAG ATATAGAGGA CAATGGGAGA AGTGAATTAT ACAATATATA ACTAGTAAAA ATTGAACCAT TAGGAGTAGC ACCACCAAG
 468 F R P G G G D M R D N W R S E L Y K Y K V V K I E P L G V A P T K
 earI/ksp632I styI alwNI
 1501 GCAATGAGAA GAGTGGTGCA GAGAGAAAA AGAGCGTGG GAATAGGAGC TGCTTCCTT GGGTCTTAG GAGCAGCAGG AAGCACTATG GCGCAGCGT
 CGTTACTCTT CTCACCCAGT CTCTCTTTT TCTCGTCACC CTTATCCTCG ACACAAGGAA CCAAGAATC CTCGTGTC TCCTGATAC CCGCGTCGCA
 501 A M R R V V Q R E K R A V G I G A V F L G A A G S T M G A A S
 haeI alwNI
 1601 CAATAACGCT GACGGTACAG GCCAGACTAT TATTGCTGG TATAGTGCAA CAGCAGAACAA ATTGCTGAG GGTATTGAG GCGCAACAGC ATCTGTTGCA
 GTTATTGCA CTGCCATGTC CCGTCTGATA ATACAGACC ATATCAGTT GTGCTTGT TARACGACT CCGATAACTC CCGTGTGTC TAGACACGT
 535 I T L T V Q A R L L L S G I V Q Q N N L L R A I E A Q Q H L L Q
 gsuI/bpmI eco8II alwNI
 1701 ACTCACAGTC TGGGGCATCA AGCAGCTCCA GCAAGAGTC CTGGCTGTCG AAGATACCT AAGGATCAA CAGCTCCTGG GGATTGGGG TTGCTCTGGA
 TGAGTGTGAG ACCCGTAGT TCGTCGAGT CCGTCTCAG GACCGACACC TTTCTATGGA TTCCCTAGTT GTCGAGGACC CCTAACCC CACGAGACCT
 568 L T V W G I K Q L Q A R V L A V E R Y L R D Q Q L L G I W G C S G
 styI bsmI xbaI
 1801 AAACCTCATTT GCACCACTC TGTCCTTGG AATGCTAGTT GGAGTATATA ATCTCTAGAT AAGATTGGG ATACATGAC CTGGATGGAG TGGGAAGAG
 TTGAGTAAA CGTGGTGGAG ACACGGAACC TTACGATCAA CCTCATTTT TAGAGATCTA TTCTAAACCC TATTCTACTG GACTACCTC ACCCTTCTC
 601 K L I C T T S V P W N A S W S N K S L D K I W D N M T W M E W E R E
 hindIII
 1901 AAATTGAGAA TTACACRAGC TTAATATACA CCTTAATTGA AGAATCGCAG AACCAACRAG AACAGACTTA TTGCAATTGG ATCAATAGCG
 TTTAACTCTT AATGTTGTCG AATTATATGT GGAATTAAT TCTTAGCGTC TTGTTGTTT TTTCTTATT TGTTCTGAAT AACCTAAC TAGTTATCCG
 635 I E N Y T S L I Y T L I E S Q N Q Q E K N K Q D L L E L D Q O A
 ssfI
 2001 AAGTTGTGG AATTGGTTTA GCATAACAAA ATGGCTGTGG TATATAAAA TATTCTAAT GATAGTTGGA GGCTTGTAG GTTAAGAAT AGTTTTCCT
 TTCACACACC TTAACCAAT CGTATTGTTT TACCGACACC ATATATTTT ATAAGTATA CTATCAACT CCGAACCATC CAAATTCTTA TCRAAAGGA
 668 S L W N W F S I T K W L W Y I K I F I M I V G G L V G L R I V F A
 scfI ppuNI
 2101 GTACTTTCTA TAGTGAATAG AGTAGGCGAG GGTACTCAC CATTATCAAT TCAGACCCGC CTCGAGGCC CGAGGGGACC CGACAGGCC AAAGGAATCG
 CATGAAGAT ATCACTTATC TCAATCCGTC CCGATGAGTG GTAAATAGTAA AGTCTGGCG GAGGTCGGG GCTCCCTGG GCTGTCGGG TTTCCTTAGC
 701 V L S I V N R V R Q G Y S P L S F Q T R L P A P R G P D R P K G I E

2201 AAGAGAAGG TGGAGAGCA GACAGGACA GATCCATTG CTTAGTGGAT GGATTCTTAG CACTTATCTG GGACGATCTA CGGAGCCTCT GCCTCTTCAG
 TTCTTCTCC ACCTCTCGT CTGTCCTGT CTAGTAAGC GATCACCTA CTAAGAATC GTGAATAGAC CTTGCTAGAT GCCTGGACA CGGAGAAGTC
 735 E E G G E Q D R D R S I R L V D G F L A L I W D D L R S L C L F S
 2301 CTACCACCGC TTGAGAGACT TACTCTTGAT TGCAACGAGG ATTCTGGAAC TTCTGGGACG CAGGGGGTGG GAAGCCCTCA AATATTGGTG GAATCTCCTA
 GATGGTGCG AACTCTCTGA ATGAGAACTA ACGTTGCTCC TAACACCTTG AAGACCCCTG GTCCCCCACC CTTGGGGAGT TTATAACCAC CTTAGAGGAT
 768 Y H R L R D L L L I A T R I V E L L G R R G W E A L K Y W N L L
 2401 CAGTATTGGA TTCAGGAACT AAAGAAATAGT GCTGTTAGCT TGTTAATGT CACAGCCATA GCAGTAGCTG AGGGACAGA TAGGGTTCTA GAAGCATTGC
 GTCATAACCT AAGTCCCTGA TTCTTATCA CGACAATCGA ACGAATTACA GTCTGGGTAT CGTCATCGAC TCCCCTGTCT ATCCCAAGAT CTTGCTAACG
 801 Q Y W I Q E L K N S A V S L L N V T A I A V A E G T D R V L E A L Q
 2501 AAAGAGCTTA TAGAGCTATT CTCCACATAC CTACAGAAT AAGACAAGGC TTGAAAGGG CTTTGCTATA A
 TTCTCGAAT ATCTCGATAA GAGGTGTATG GATGTTCTTA TTCGTTCGG AACCTTTCCC GAAACGATAT T
 835 R A Y R A I L H I P T R I R Q G L E R A L L O

Table 3 illustrates the amino acid sequences for the MN, GNE₈, and GNE₁₆ gp120 proteins. The regions of the sequences having identical amino acid sequences are enclosed in boxes.

5

TABLE 3

gp160.8.24	1	M	I	V	K	G	I	R	K	N	C	O	H	L	W	R	W	G	T	M	L	L	G	M	L	M	I	C	S	A	A	E	K	L	W	V	T	V	Y	Y	G	V	P	V	W	K	E	A	T	T	T	
gp160.SF.16.2	1	M	R	V	K	G	I	R	R	N	Y	O	H	L	W	R	W	G	T	M	L	L	G	I	L	M	I	C	S	A	A	G	K	L	W	V	T	V	Y	Y	G	V	P	V	W	K	E	T	T	T	T	
gp160.SF.16.7	1	M	R	V	K	R	I	R	R	N	Y	O	H	L	W	K	W	G	T	M	L	L	G	M	L	M	I	C	S	A	A	G	K	L	W	V	T	V	Y	Y	G	V	P	V	W	K	E	T	T	T	T	
gp160.8.24	51	L	F	C	A	S	D	A	K	A	Y	D	T	E	I	H	N	V	W	A	T	H	A	C	V	P	T	D	P	N	P	O	E	I	G	L	E	N	V	T	E	N	F	N	M	W	K	N	N	M	V	
gp160.SF.16.2	51	L	F	C	A	S	D	A	K	A	Y	D	T	E	I	H	N	V	W	A	T	H	A	C	V	P	T	D	P	N	P	O	E	V	V	L	E	N	V	T	E	N	F	N	M	W	K	N	N	M	V	
gp160.SF.16.7	51	L	F	C	A	S	D	A	K	A	Y	D	T	E	I	H	N	V	W	A	T	H	A	C	V	P	T	D	P	N	P	O	E	V	V	L	E	N	V	T	E	N	F	N	M	W	K	N	N	M	V	
gp160.8.24	101	E	O	M	H	E	D	I	I	S	L	W	D	O	S	L	K	P	C	V	K	L	T	P	L	C	V	T	L	N	C	T	D	L	K	N	A	T	N	T	T	S	S	S	W	G	K	M	E	R	G	
gp160.SF.16.2	101	E	O	M	H	E	D	I	I	S	L	W	D	O	S	L	K	P	C	V	K	L	T	P	L	C	V	T	L	N	C	T	D	A	G	N	T	T	N	T	N	S	S	S	R	E	K	L	E	K	G	
gp160.SF.16.7	101	E	O	M	H	E	D	I	I	S	L	W	D	O	S	L	K	P	C	V	K	L	T	P	L	C	V	T	L	N	C	T	D	A	G	N	T	T	N	T	N	S	S	S	G	E	K	L	E	K	G	
gp160.8.24	151	E	I	K	N	C	S	F	N	I	T	T	S	I	R	D	K	M	K	N	E	V	A	L	F	Y	K	L	D	V	P	I	D	N	D	N	T	S	Y	R	L	I	S								
gp160.SF.16.2	151	E	I	K	N	C	S	F	N	I	T	T	S	V	R	D	K	M	O	K	E	T	A	L	F	N	K	L	D	I	V	P	I	D	D	D	R	N	S	T	R	N	S	T	N	Y	R	L	I	S		
gp160.SF.16.7	151	E	I	K	N	C	S	F	N	I	T	T	S	M	R	D	K	M	O	R	E	T	A	L	F	N	K	L	D	I	V	P	I	D	D	D	R	N	S	T	R	N	S	T	N	Y	R	L	I	S		
gp160.8.24	194	C	N	T	S	V	I	T	O	A	C	P	K	V	S	F	E	P	I	P	I	H	Y	C	A	P	A	G	F	A	L	L	K	C	R	D	K	K	F	N	G	T	G	P	C	T	N	V	S	T	V	
gp160.SF.16.2	201	C	N	T	S	V	I	T	O	A	C	P	K	V	S	F	E	P	I	P	I	H	F	C	T	P	A	G	F	A	L	L	K	C	N	N	K	T	F	N	G	S	G	P	C	K	N	V	S	T	V	
gp160.SF.16.7	201	C	N	T	S	V	I	T	O	A	C	P	K	V	S	F	E	P	I	P	I	H	F	C	T	P	A	G	F	A	L	L	K	C	N	N	E	T	F	N	G	S	G	P	C	K	N	V	S	T	V	
gp160.8.24	244	O	C	T	H	G	I	R	P	V	V	S	T	O	L	L	N	G	S	L	A	E	E	E	V	V	I	R	S	A	N	F	S	D	N	A	K	T	I	I	V	O	L	N	E	S	V	E	I	N		
gp160.SF.16.2	251	O	C	T	H	G	I	R	P	V	V	S	T	O	L	L	N	G	S	L	A	E	G	E	V	V	I	R	S	E	N	F	T	N	N	A	K	T	I	I	V	O	L	T	E	P	V	K	I	N		
gp160.SF.16.7	251	L	C	T	H	G	I	R	P	V	V	S	T	O	L	L	N	G	S	L	A	E	G	E	V	V	I	R	S	E	N	F	T	N	N	A	K	T	I	I	V	O	L	K	E	P	V	K	I	N		
gp160.8.24	294	C	T	R	P	N	N	N	T	R	S	I	H	I	G	P	G	R	A	F	Y	A	T	G	E	I	I	G	D	I	R	O	A	H	C	N	L	S	S	T	K	W	N	T	L	K	O	I	V			
gp160.SF.16.2	301	C	T	R	P	N	N	N	T	R	K	S	I	P	I	G	P	G	R	A	F	Y	A	T	G	D	I	I	G	N	I	R	O	A	H	C	N	L	S	R	T	D	W	N	N	T	L	G	O	I	V	
gp160.SF.16.7	301	C	T	R	P	N	N	N	T	R	K	S	I	P	I	G	P	G	R	A	F	Y	A	T	G	D	I	I	G	N	I	R	O	A	H	C	N	L	S	R	T	D	W	N	N	T	L	R	O	I	A	
gp160.8.24	344	T	K	L	R	E	H	F	-	N	K	T	I	I	F	N	H	S	S	G	G	D	P	E	I	V	M	H	S	F	N	C	R	G	E	F	F	Y	C	N	T	T	P	L	F	N	S	T	W	N	Y	
gp160.SF.16.2	351	E	K	L	R	E	O	F	G	N	K	T	I	I	F	N	H	S	S	G	G	D	P	E	I	V	M	H	S	F	N	C	R	G	E	F	F	Y	C	N	T	T	Q	L	F	D	S	T	W	D	N	
gp160.SF.16.7	351	E	K	L	R	K	O	F	G	N	K	T	I	I	F	N	H	S	S	G	G	D	P	E	I	V	M	H	S	F	N	C	R	G	E	F	F	Y	C	D	T	Q	L	F	N	S	T	W	N	A		
gp160.8.24	393	T	Y	T	W	N	N	T	E	G	S	N	O	T	G	R	N	I	T	L	Q	C	R	I	K	O	I	I	N	M	W	Q	E	V	G	K	A	M	Y	A	P	P	I	R	G	O	I	R	C	S	S	
gp160.SF.16.2	401	T	K	V	-	-	S	N	G	T	S	T	E	E	N	S	T	I	T	L	P	C	R	I	K	O	I	V	N	M	W	Q	E	V	G	K	A	M	Y	A	P	P	I	R	G	O	I	R	C	S	S	
gp160.SF.16.7	401	N	N	T	-	-	E	R	-	N	S	T	K	E	N	S	T	I	T	L	P	C	R	I	K	O	I	V	N	M	W	Q	E	V	G	K	A	M	Y	A	P	P	I	R	G	O	I	R	C	S	S	
gp160.8.24	443	N	I	T	G	L	L	L	T	R	O	G	G	-	N	N	S	E	T	E	I	I	F	R	P	G	G	G	D	M	R	D	N	W	R	S	E	L	Y	K	Y	K	V	V	K	I	E	P	L	G	V	A
gp160.SF.16.2	449	N	I	T	G	L	L	L	T	R	O	G	G	S	N	N	S	M	N	E	T	I	F	R	P	G	G	G	D	M	R	D	N	W	R	S	E	L	Y	K	Y	K	V	V	K	I	E	P	L	G	V	A
gp160.SF.16.7	448	N	I	T	G	L	L	L	T	R	O	G	G	S	S	N	S	M	N	E	T	I	F	R	P	G	G	G	D	M	R	D	N	W	R	S	E	L	Y	K	Y	K	V	V	K	I	E	P	L	G	V	A
gp160.8.24	492	P	T	K	A	K	R	R	V	M	O	R	E	K	R	A	V	G	I	G	A	V	F	L	G	F	L	G	A	A	G	S	T	M	G	A	A	S	V	T	L	T	V	Q	A	R	L	L	S	G		
gp160.SF.16.2	499	P	T	K	A	K	R	R	V	V	O	R	E	K	R	A	V	G	I	G	A	V	F	L	G	F	L	G	A	A	G	S	T	M	G	A	A	S	I	T	L	T	V	Q	A	R	L	L	S	G		
gp160.SF.16.7	498	P	T	K	A	M	R	R	V	V	O	R	E	K	R	A	V	G	I	G	A	V	F	L	G	F	L	G	A	A	G	S	T	M	G	A	A	S	I	T	L	T	V	Q	A	R	L	L	S	G		
gp160.8.24	542	I	V	Q	O	O	N	N	L	L	R	A	I	E	A	O	H	L	L	O	L	T	I	V	W	G	I	K	O	L	Q	A	R	V	L	A	V	E	R	Y	L	K	D	O	Q	L	L	G	I	W	G	
gp160.SF.16.2	549	I	V	Q	O	O	N	N	L	L	R	A	I	E	A	O	H	L	L	O	L	T	I	V	W	G	I	K	O	L	Q	A	R	V	L	A	V	E	R	Y	L	R	D	O	Q	L	L	G	I	W	G	
gp160.SF.16.7	548	I	V	Q	O	O	N	N	L	L	R	A	I	E	A	O	H	L	L	O	L	T	I	V	W	G	I	K	O	L	Q	A	R	V	L	A	V	E	R	Y	L	R	D	O	Q	L	L	G	I	W	G	

gp160.8.24	592	CSGKLICTTA	VPWNASWSNKS	LDKIWDNMTWMEW	REIDNYTSLIYS	SLIE
gp160.SF.16.2	599	CSGKLICTTS	VPWNASWSNKS	LDKIWDNMTWMEW	REIDNYTSLIYT	LIE
gp160.SF.16.7	598	CSGKLICTTS	VPWNASWSNKS	LDKIWDNMTWMEW	REIDNYTSLIYT	LIE

gp160.8.24	642	ESONQOEKNEQ	ELLELDK	WASLWNWF	DI	TKWLWYIKIFIMIVGG	LVGLRI
gp160.SF.16.2	649	ESONQOEKNEQ	ELLELDK	WASLWNWF	SI	TKWLWYIKIFIMIVGG	LVGLRI
gp160.SF.16.7	648	ESONQOEKNEQ	ELLELDK	WASLWNWF	SI	TKWLWYIKIFIMIVGG	LVGLRI

stop

gp160.8.24	692	VFTVLSIVNRVR	KGYSPLSFQTH	LPAPRGL	DRPEGT	EEEGGERDRDRSS	IR
gp160.SF.16.2	699	VFAVLSIVNRVR	QGYSPISFQTR	LPAPRRP	DRPEGIE	EEEGGEQGRDRS	IR
gp160.SF.16.7	698	VFAVLSIVNRVR	QGYSPISFQTR	LPAPRGP	DRPKGIE	EEEGGEQGRDRS	IR

gp160.8.24	742	LDGFLAIVW	DLRSLCLFSYHRL	RDLLIAA	RIVELLGRRGWEAL	KYWW
gp160.SF.16.2	749	LDGFLALIW	DLRSLCLFSYHRL	RDLLIAT	RIVELLGRRGWEAL	KYWW
gp160.SF.16.7	748	LDGFLALIW	DLRSLCLFSYHRL	RDLLIAT	RIVELLGRRGWEAL	KYWW

gp160.8.24	792	NLLOYWIOELKNSAVSLLN	ATAIAVAEGTDRV	IEIV	ORAYRAILHIPTRI
gp160.SF.16.2	799	NLLOYWIOELKNSAVSLLN	VTAIAVAEGTDRV	LEV	ORAYRAILHIPTRI
gp160.SF.16.7	798	NLLOYWIOELKNSAVSLLN	VTAIAVAEGTDRV	LEAL	ORAYRAILHIPTRI

gp160.8.24	842	ROGLERALL
gp160.SF.16.2	849	ROGLERALL
gp160.SF.16.7	848	ROGLERALL

Nucleic acid sequences encoding gp120 from GNE₈ and GNE₁₆ capable of expressing gp120 can be prepared by conventional means. The nucleotide sequence can be synthesized. Alternatively, another HIV nucleic acid sequence encoding gp120 can be used as a backbone and altered at any differing residues by site directed mutagenesis as described in detail in Example 1.

In a preferred embodiment, the nucleotide sequence is present in an expression construct containing DNA encoding gp120 under the transcriptional and translational control of a promoter for expression of the encoded protein. The promoter can be a eukaryotic promoter for expression in a mammalian cell. In cases where one wishes to expand the promoter or produce gp120 in a prokaryotic host, the promoter can be a prokaryotic promoter. Usually a strong promoter is employed to provide high level transcription and expression.

The expression construct can be part of a vector capable of stable extrachromosomal maintenance in an appropriate cellular host or may be integrated into host genomes. Normally, markers are provided with the expression construct which allow for selection of a host containing the construct. The marker can be on the same or a different DNA molecule, desirably, the same DNA molecule.

The expression construct can be joined to a replication system recognized by the intended host cell. Various replication systems include viral replication systems such as retroviruses, simian virus, bovine papilloma virus, or the like. In addition, the construct may be joined to an amplifiable gene, e.g. DHFR gene, so that multiple copies of the gp120 DNA can be made. Introduction of the construct into the host will vary depending on the construct and can be

achieved by any convenient means. A wide variety of prokaryotic and eukaryotic hosts can be employed for expression of the proteins.

Preferably, the gp120 is expressed in mammalian cells that provide the same glycosylation and disulfide bonds as in native gp120. Expression of gp120 and fragments of gp120 in mammalian cells as fusion proteins incorporating N-terminal sequences of Herpes Simplex Virus Type 1 (HSV-1) glycoprotein D (gD-1) is described in Lasky, L. A. et al., 1986 (Neutralization of the AIDS retrovirus by antibodies to a recombinant envelope glycoprotein) Science 233: 209-212 and Haffar, O.K. et al., 1991 (The cytoplasmic tail of HIV-1 gp160 contains regions that associate with cellular membranes.) Virol. 180:439-441, respectively. A preferred method for expressing gp120 is described in Example 3. In the example, a heterologous signal sequence was used for convenient expression of the protein. However, the protein can also be expressed using the native signal sequence.

An isolated, purified GNE₈-gp120 and GNE₁₆-gp120 having the amino acid sequence illustrated in Tables 1-3 can be produced by conventional methods. For example, the proteins can be chemically synthesized. In a preferred embodiment, the proteins are expressed in mammalian cells using an expression construct of this invention. The expressed proteins can be purified by conventional means. A preferred purification procedure is described in Example 3.

30

gp120 Fragments

The present invention also provides gp120 fragments that are suitable for use in inducing antibodies for use in serotyping or in a vaccine formulation. A truncated gp120 sequence as used herein is a fragment of gp120 that is free from a portion of

35

the intact gp120 sequence beginning at either the amino or carboxy terminus of gp120. A truncated gp120 sequence of this invention is free from the C5 domain. The C5 domain of gp120 is a major immunogenic site of the molecule. However, antibodies to the region do not neutralize virus. Therefore, elimination of this portion of gp120 from immunogens used to induce antibodies for serotyping is advantageous.

In another embodiment, the truncated gp120 sequence is additionally free from the carboxy terminus region through about amino acid residue 453 of the gp120 V5 domain. The portion of the V5 domain remaining in the sequence provides a convenient restriction site for preparation of expression constructs. However, a truncated gp120 sequence that is free from the entire gp120 V5 domain is also suitable for use in inducing antibodies.

In addition, portions of the carboxy terminus of gp120 can also be eliminated from the truncated gp120 sequence. The truncated gp120 sequence can additionally be free from the gp120 signal sequence. The truncated gp120 sequence can be free from the carboxy terminus through amino acid residue 111 of the gp120 C1 domain, eliminating most of the C1 domain but preserving a convenient restriction site. However, the portion of the C1 domain through the cysteine residue that forms a disulfide bond can additionally be removed, so that the truncated gp120 sequence is free from the carboxy terminus through amino acid residue 117 of the gp120 C1 domain. Alternatively, the truncated gp120 sequence can be free from the amino terminus of gp120 through residue 111 of the C1 domain, preserving the V2 disulfide bond. In a preferred embodiment, the truncated gp120 sequence is free from the amino terminus of gp120 through residue 111 of the

C1 domain and residue 453 through the carb xy terminus of gp120.

The truncated gp120 sequences can be produced by recombinant engineering, as described previously.

- 5 Conveniently, DNA encoding the truncated gp120 sequence is joined to a heterologous DNA sequence encoding a signal sequence.

Serotyping method

- 10 The present invention also provides an improved serotyping method for HIV strains. The method comprises determining the serotypes of the V2, V3, and C4 domains of gp120.

- HIV isolates can be serotyped by conventional
15 immunoassay methods employing antibodies to the neutralizing epitopes in the V2, V3, and C4 domains for various strains of HIV. Preparation of the antibodies is described hereinbefore. The antibody affinity required for serotyping HIV using a particular
20 immunoassay method does not differ from that required to detect other polypeptide analytes. The antibody composition can be polyclonal or monoclonal, preferably monoclonal.

- A number of different types of immunoassays are
25 well known using a variety of protocols and labels. The assay conditions and reagents may be any of a variety found in the prior art. The assay may be heterogeneous or homogeneous. Conveniently, an HIV isolate is adsorbed to a solid phase and detected with
30 antibody specific for one strain of neutralizing epitope for each neutralizing epitope in the V2, V3, and C4 domain. Alternatively, supernatant or lysate from the cultured isolate which contains gp120 can be adsorbed to the solid phase. The virus or gp120 can be
35 adsorbed by many well known non-specific binding methods. Alternatively, an anti-gp120 antibody,

preferably directed to the carboxy terminus of gp120. can be used to affix gp120 to the solid phase. A gp120 capture antibody and sandwich ELISA assay for gp120 neutralizing epitopes is described by Moore, *AIDS Res. Hum. Retroviruses* 9:209-219 (1993). Binding between the antibodies and sample can be determined in a number of ways. Complex formation can be determined by use of soluble antibodies specific for the anti-gp120 antibody. The soluble antibodies can be labeled directly or can be detected using labeled second antibodies specific for the species of the soluble antibodies. Various labels include radionuclides, enzymes, fluorescers, colloidal metals or the like. Conveniently, the anti-gp120 antibodies will be labeled directly, conveniently with an enzyme.

Alternatively, other methods for determining the neutralizing epitopes can be used. For example, fluorescent-labeled antibodies for a neutralizing epitope can be combined with cells infected by the strain of HIV to be serotyped and analyzed by fluorescence activated cell sorting.

The serotype of the HIV isolate includes the strain of the neutralizing epitopes for the V2, V3, and C4 domains.

It is understood that the application of the teachings of the present invention to a specific problem or situation will be within the capabilities of one having ordinary skill in the art in light of the teachings contained herein. Examples of the products of the present invention and representative processes for their isolation, use, and manufacture appear below, but should not be construed to limit the invention. All literature citations herein are expressly incorporated by reference.

EXAMPLE 1**Identification of C4 Neutralizing Epitopes**

The following reagents and methods were used in the studies described herein.

5

gp120 sequences and nomenclature. Amino acid residues are designated using the standard single letter code. The location of amino acids within the gp120 protein is specified using the initiator methionine residue as position 1. The designation LAI is used to describe the virus isolate from which the HIV-1_{BH10}, HIV-1_{MB}, HIV-1_{BRU}, HIV-1_{HXB2}, HIV-1_{HXB3} and HIV-1_{HXB10} substrains (molecular clones) of HIV-1 were obtained. The sequence of gp120 from IIIB substrain of HIV-1_{LAI} is that determined by Muesing et al. (30).

15

The sequence of gp120 from MN strain of HIV-1 is given with reference to the MNgp120 clone (MN_{GNE}). The sequence of this clone differs by approximately 2% from that of the MN₁₉₈₄ clone described by Gurgo et al. (13). The sequences of gp120 from the NY-5, JRcsf, Z6, Z321, and HXB2 strains of HIV-1 are those listed by Myers et al. (32) except where noted otherwise. The sequence of the Thai isolate A244 is that provided by McCutchan et al. (24). The variable (V) domains and conserved (C) domains of gp120 are specified according to the nomenclature of Modrow et al. (28).

20

25

Monoclonal antibody production and screening assays. Hybridomas producing monoclonal antibodies to MN-rgp120 (recombinantly produced gp120 from the MN strain of HIV) (3) were prepared and screened for CD4 blocking activity as described previously (7, 33). The binding of monoclonal antibodies to MN-rgp120 and to rgp120s from the IIIB, NY-5, Z6, Z321, JRcsf, and A244 strains of HIV-1 was assessed by enzyme linked

30

35

immunoabsorbant assays (ELISA) as described previously (33).

Virus binding and neutralization assays. The ability of monoclonal antibodies to neutralize HIV-1 infectivity *in vitro* was assessed in a colorimetric MT-2 cell cytotoxicity assay similar to that described previously (35). MT-2 cells and H9/HTLV-III_{MN} cells were obtained through the AIDS Research and Reference Reagent Program, Division of AIDS, NIAID, NIH: contributed by Drs. Douglas Richman and Robert Gallo, respectively. Briefly, serial dilutions of antibody or serum were prepared in 50 μ l volumes of complete and then 50 μ l of a prediluted HIV-1 stock was added to each well. After incubation for 1 hr at 4°C, 100 μ l of a 4×10^5 MT-2 cell/ml suspension was added. After incubation of the plates for 5 days at 37°C in 5% CO₂, viable cells were measured using metabolic conversion of the formazan MTT dye. Each well received 20 μ l of a 5 mg/ml MTT solution in PBS.

After a 4 hr incubation at 37°C, the dye precipitate was dissolved by removing 100 μ l of the cell supernatant, adding 130 μ l of 10% Triton X-100 in acid isopropanol, then pipeting until the precipitate was dissolved. The optical density of the wells was determined at 540 nm. The percentage inhibition was calculated using the formula:

$$\frac{1 - (\text{virus control} - \text{experimental})}{(\text{virus control} - \text{medium control})}$$

Cell surface staining of HIV-1 infected cells with monoclonal antibodies. H9 cells (2×10^5) chronically infected with the IIIB, HXB2, HXB3, and HX10 substrains of HIV-1_{LAI} or with HIV-1_{MN} were incubated for 30 min at room temperature with monoclonal antibodies (10 μ g per ml) in 100 μ l of RPMI 1640 cell culture media

containing 1% FCS. Cells were washed and then incubated with 20 μ g per ml of fluorescein-conjugated, affinity-purified, goat antibody to mouse IgG (Fab'), (Cappel, West Chester, PA) for 30 min. Cells were washed, fixed with 1% paraformaldehyde and the bound antibody was quantitated by flow cytometry using a FACSCAN (Becton-Dickenson, Fullerton, CA).

Fluorescence data was expressed as percentage of fluorescent cells compared to the fluorescence obtained with the second antibody alone. Fluorescence was measured as the mean intensity of the cells expressed as mean channel number plotted on a log scale.

Fragmentation of the MN-rgp120 gene. Fragments of the MN-rgp120 gene were generated using the polymerase chain reaction (PCR) (17). Briefly, forward 30-mer oligonucleotide DNA primers incorporating a Xho 1 site, and reverse 36-mer oligonucleotide DNA primers containing a stop codon followed by a Xba 1 site were synthesized and used for the polymerase chain reactions. Thirty cycles of the PCR reaction were performed using 0.3 μ g of a plasmid containing the gene for gp120 from the MN strain of HIV-1 (pRKMN. D533) and 0.04 nM of a designated primers. The PCR reaction buffer consisted of 0.1 M Tris buffer (pH 8.4), 50 mM KCl, 0.2 mM 4dNTP (Pharmacia, Piscataway, NJ), 0.15 M MgCl₂, and 0.5 Unit of Taq Polymerase (Perkin-Elmer Cetus, Norwalk, CN) and a typical PCR cycle consisted of a 60 second denaturation step at 94°C, followed by a 45 second annealing step at 55° C, and then an extension step at 72° C for 45 seconds.

Following the PCR amplification, the PCR products were purified by phenol and chloroform extraction, and then ethanol precipitated. The purified products were then digested with the restriction endonucleases Xho1 and Xba1. The resulting PCR products were gel purified

using 1% agarose (SEAKEM, FMC Bioproducts, Rockland, ME) or 5% polyacrylamide gel electrophoresis (PAGE) and then isolated by electroelution.

5 **Site directed mutagenesis of the MN-rgp120 C4 domain.** A recombinant PCR technique (15) was utilized to introduce single amino acid substitutions at selected sites into a 600 bp Bgl II fragment of MN-rgp120 that contained the C4 domain. This method
10 entailed the PCR amplification of overlapping regions of the C4 domain of gp120 using primers that incorporated the desired nucleotide changes. The resultant PCR products were then annealed and PCR
15 amplified to generate the final product. For these reactions 18-mer "outside" primers encoding the wild type sequence (Bgl II sites) were amplified with 36-mer "inside" primers that contained the alanine or glutamic acid residue changes. The first PCR reaction included
20 1X of the Vent polymerase buffer (New England Biolabs, Beverly, MA), 0.2 mM of 4dNTP (Pharmacia, Piscataway, N.J.), 0.04 nM of each synthetic oligonucleotide, 0.3 µg of linearized plasmid, pRKMN.D533, which contained the MN-rgp120 gene. Thirty PCR cycles were performed consisting of the following sequence of steps: 45
25 seconds of denaturation at 94°C, 45 second of annealing at 55°C and 45 seconds of extension at 72°C. Following PCR amplification, the product pairs were gel purified using a 1% solution of low melt agarose (SeaPlaque, FMC Bioproducts, Rockland, ME).
30 The agarose containing PCR product was melted at 65°C and combined with the PCR product of the overlapping pair and equilibrated to 37°C. Added to this (20 µl) was 10 µl of 10X Vent Polymerase buffer, 10 µl of 2 mM 4dNTP, 0.04 nM each of the "outside" wild
35 type 18 mer oligonucleotides, 57 µl of H₂O and 1 unit of

Vent Polymerase. Thirty PCR cycles were performed as previously above.

The resulting PCR products were purified and digested with the Bgl II endonuclease. The digested PCR product was then ligated into the mammalian cell expression vector pRKMN.D533, which had been digested with Bgl II allowing for the removal of a 600 bp fragment. Colonies containing the correct insertion were identified and Sequenase 2.0 supercoil sequencing was employed to check for fidelity and the incorporation of the desired mutation.

Expression of gp120 fragments in mammalian cells. Fragments of the MN and IIIB gp120 were expressed in mammalian cells as fusion proteins incorporating N-terminal sequences of Herpes Simplex Virus Type 1 (HSV-1) glycoprotein D (gD-1) as described previously (14, 22). Briefly, isolated DNA fragments generated by the PCR reaction were ligated into a plasmid (pRK.gD-1) designed to fuse the gp120 fragments, in frame, to the 5' sequences of the glycoprotein D (gD) gene of Type 1 Herpes Simplex Virus (gD-1) and the 3' end to translational stop codons. The fragment of the gD-1 gene encoded the signal sequence and 25 amino acids of the mature form of HSV-1 protein. To allow for expression in mammalian cells, chimeric genes fragments were cloned into the pRK5 expression plasmid (8) that contained a polylinker with cloning sites and translational stop codons located between a cytomegalovirus promotor and a simian virus 40 virus polyadenylation site.

The resulting plasmids were transfected into the 293s embryonic human kidney cell line (12) using a calcium phosphate technique (11). Growth conditioned cell culture media was collected 48 hr after transfection, and the soluble proteins were detected by

ELISA or by specific radioimmunoprecipitation where metabolically labeled proteins from cell culture supernatants were resolved by sodium dodecyl sulfate polyacrylamide gel electrophoresis (PAGE) and
5 visualized by autoradiography as described previously (1, 18).

Radioimmunoprecipitation of MN-rgp120 mutants.
Plasmids directing the expression of the MN-rgp120 C4
10 domain mutants were transfected into 293s cells as described above. Twenty four hours following the transfection, the cells were metabolically labeled with [³⁵S]-labeled methionine or cysteine as described previously (1). The labeled cell culture supernatants
15 were then harvested and 0.5 ml aliquots were reacted with 1-5 µg of the monoclonal antibody or with 2 µl of the polyclonal rabbit antisera to MN-rgp120 and immunoprecipitated with Pansorbin (CalBiochem, La Jolla, CA) as described previously (1). The resulting
20 Pansorbin complex was pelleted by centrifugation, washed twice with a solution containing PBS, 1% NP-40 and 0.05% SDS and then boiled in a PAGE sample buffer containing 1% 2-mercaptoethanol. The processed samples were then analyzed by SDS-PAGE and visualized by
25 autoradiography (1, 18).

Assays to measure the binding of monoclonal antibodies to mutagenized MN-rgp120 polypeptides. An ELISA was developed to screen for reactivity of
30 MN-rgp120 fragments and mutant proteins with various monoclonal antibodies. In this assay, 96 well microtiter dishes (Maxisorp, Nunc, Roskilde, Denmark) were coated overnight with mouse monoclonal antibody (5B6) to gD-1, at a concentration of 2.0 µg/ml in
35 phosphate buffered saline (PBS). The plates were blocked in a PBS solution containing 0.5% bovin serum

albumin (PBSA) and then incubated with growth conditioned cell culture medium from transfected cells expressing the recombinant gp120 variants for 2 hr at room temperature. The plates were washed three times in PBS containing 0.05% Tween 20 and then incubated with the purified, HRP-conjugated monoclonal antibodies. Following a 1 hr incubation, the plates were washed three times and developed with the colorimetric substrate, o-phenylenediamine (Sigma, St. Louis, MO).

The optical densities in each well were then read in a microtiter plate reading spectrophotometer at 492 nm. Each cell culture supernatant containing fragments or mutated rgp120s was normalized for expression based on the titering of its reactivity to the V3 binding monoclonal antibody 1034 or to a rabbit polyclonal antiserum to MN-rgp120. Data from these experiments were expressed as a ratio of the optical densities obtained with the CD4 blocking monoclonal antibodies binding to the fragments or MN-rgp120 mutants compared with the full length wild type rgp120s.

To normalize for different concentrations of MN-rgp120-derived protein in the cell culture supernatants, the binding of the CD4 blocking monoclonal antibodies to each preparation was compared to that of an HRP-conjugated monoclonal antibody to the V3 domain of MN-rgp120 (1034). Data from these experiments were expressed as a ratio of the optical densities obtained with the CD4 blocking monoclonal antibodies to the HRP conjugated V3 reactive monoclonal antibody.

CD4 binding assays. The ability of monoclonal antibodies to inhibit the binding of MN-rgp120 to recombinant soluble CD4 (rsCD4) was determined in a

solid phase radioimmunoassay similar to that described previously (33). The effect of single amino acid substitutions on the binding of MN-rgp120 mutants to CD4 was determined in a co-immunoprecipitation assay similar to that described previously (21). Briefly, 293 cells were metabolically labeled with ^{35}S -methionine 24 hr after transfection with plasmids expressing MN-rgp120 variants. Growth conditioned cell culture medium (0.5 ml) was then incubated with 5.0 μg of recombinant sCD4 for 90 minutes at room temperature. Following this incubation, 5.0 μg of an anti-CD4 monoclonal antibody (465), known to bind to an epitope remote from the gp120 binding site, was added and allowed to incubate another 90 minutes at room temperature.

The gp120-CD4-antibody complexes were precipitated with Pansorbin that had been washed with PBS, preabsorbed with 0.1% bovine serum albumin and then bound with 50 μg of an affinity purified rabbit anti-mouse IgG (Cappel, West Chester, PA). The pellet was washed twice with PBS 1% NP-40, 0.05% SDS, and then boiled in beta mercaptoethanol containing SDS-PAGE sample buffer. The immunoprecipitation products were resolved by SDS PAGE and visualized by autoradiography as described previously (1, 21).

Antibody affinity measurements. Anti-gp120 antibodies were iodinated with $\text{Na } ^{125}\text{I}$ with iodogen (Pierce, Rockford, IL). Briefly, 50 μg of antibody in PBS was placed in 1.5 ml polypropylene microcentrifuge tubes coated with 50 μg of Iodogen. Two millicuries of carrier free $\text{Na}[^{125}\text{I}]$ was added. After 15 min., free ^{125}I was separated from the labeled protein by chromatography on a PD-10 column (Pierce, Rockford, IL) pre-equilibrated in PBS containing 0.5% gelatin.

Antibody concentrations following iodination were determined by ELISA to calculate specific activities.

For binding assays, 96-well microtiter plates were coated with 100 μ l/well of a 10 μ g/ml solution of MN-rgp120 or IIIBrgp120 in 0.1 M bicarbonate buffer, pH 9.6 and incubated for 2 hr at room temperature or overnight at 4°C. To prevent non-specific binding, plates were blocked for 1-2 hr at room temperature with 200 μ l/well of a gelatin solution consisting of PBS containing 0.5% (wt/vol) gelatin and 0.02% sodium azide. Unlabeled anti-gp120 monoclonal antibody (0 to 400 nM) was titrated (in duplicate) in situ and radiolabeled antibody was added to each well at a concentration of 0.5 nM.

After a 1-2 hr incubation at room temperature, the plate was washed 10x with the PBS/0.5% gelatin/0.02% azide buffer to remove free antibody. The antibody-gp120 complexes were solubilized with 0.1 N NaOH/0.1% SDS solution and counted in a gamma counter. The data were analyzed by the method of Scatchard (40) using the Ligand analytical software program (31). K_d values reported represent the means of four independent determinations.

RESULTS

Characterization of monoclonal antibodies to MN-rgp120 that block CD4 binding. Monoclonal antibodies prepared from mice immunized with MN-rgp120 (3, 33), were screened for the ability to bind to MN-rgp120 coated microtiter dishes by ELISA as described previously (33). Of the thirty five clones obtained, seven were identified (1024, 1093, 1096, 1097, 1110, 1112, and 1127) that were able to inhibit the binding of MN-rgp120 to recombinant CD4 in ELISA (Figure 1) or solid phase or cell surface radioimmunoassays (21, 33). Previous studies have shown that two distinct classes

of CD4 blocking monoclonal antibodies occur: those that bind to conformation dependent (discontinuous) epitopes (16, 26, 33, 35, 45) and those that bind to conformation independent (sequential) epitopes (4, 7, 21, 33, 43).

To distinguish between these two alternatives, the binding of the monoclonal antibodies to denatured (reduced and carboxymethylated) MN-rgp120 (RCM-gp120) was measured by ELISA as described previously (33). As illustrated in Table 4, below, it was found that all of the CD4 blocking monoclonal antibodies reacted with the chemically denatured protein; indicating that they all recognized conformation independent (sequential) epitopes.

Table 4
Properties of monoclonal antibodies to MN-rgp120

5	MAb	CD4 Inhi- bitors	HIV-1 mn Neutral- ization	HIV-1 mn V3	CM- rgp120	C4 Domain peptides	rg120 cross reactivity
	1024	+	+	-	+	-	2
	1093	+	+	-	+	-	2
10	1096	+	+	-	+	-	2
	1097	+	+	-	+	-	2
	1110	+	+	-	+	-	2
	1112	+	+	-	+	-	2
	1127	+	+	-	+	-	2
15	1026	-	+	+	+	-	1,2,3,4,6
	1092	-	-	-	+	-	1,2,3,4,5
	1126	-	-	-	+	-	1,2,3,5,7
	1086	-	-	-	+	-	2
	13H8	+	-	-	+	1,3	1,2,3,4,5,6,7
20							

rgp120 cross reactivity: 1, IIIB-rgp120; 2, MN-rgp120, 3, NYS-rgp120; 4, JrCSF-rgp120; 5, Z6-rgp120; 6, Z321-rgp120; 7, A244-rgp120

25

C4 domain peptides:

- 1, FINMWQEVGKAMYAPPIS (SEQ. ID. NO. 24);
- 2, MWQEVGKAMYAP (SEQ. ID. NO. 25);
- 3, GKAMYAPPIKGQIR (SEQ. ID. NO. 26)

30

The cross reactivity of these monoclonal antibodies was assessed by ELISA as described previously (33). In these experiments, the ability of the monoclonal antibodies to bind to a panel of seven rgp120s, prepared from the IIIB, MN, Z6, Z321, NY-5, A244, and JRCSF isolates of HIV-1, was measured by ELISA (33). It was found that all of the CD4 blocking monoclonal antibodies were strain specific and bound only to gp120 from the MN strain of HIV-1 (Table 4).

40 However, other antibodies from the same fusion

(1026, 1092, and 1126) exhibited much broader cross reactivity (Table 4, Figure 2), as did a CD4 blocking monoclonal antibody to IIIB-rgp120 (13H8) described previously (33).

5 Further studies were performed to characterize the neutralizing activity of the antibodies to MN-rgp120. In these studies, monoclonal antibodies were incubated with cell free virus (HIV-1_{MN}), and the resulting mixture was then used to infect MT-2 cells in
10 microtiter plates. After 5 days, the plates were developed by addition of the colorimetric dye, MTT, and cell viability was measured spectrophotometrically. It was found (Table 4, Figure 2) that all of the CD4 blocking monoclonal antibodies were able to inhibit
15 viral infectivity. However the potency of the monoclonal antibodies varied considerably with some monoclonal antibodies (eg. 1024) able to inhibit infection at very low concentrations (IC₅₀ of 0.08 µg per ml) whereas other monoclonal antibodies (eg. 1112)
20 required much higher concentrations (IC₅₀ of 30 µg per ml). In control experiments two monoclonal antibodies to MN-rgp120 from the same fusion (eg. 1086, 1092) were ineffective, whereas the 1026 monoclonal antibody exhibited potent neutralizing activity. Similarly,
25 monoclonal antibodies to the V3 domain of IIIB-rgp120 (10F6, 11G5) known to neutralize the infectivity HIV-1_{III} (33), were unable to neutralize the HIV-1_{MN} virus.

Binding studies using synthetic peptides were then performed to further localize the epitopes recognized
30 by these monoclonal antibodies as described previously (33). When a peptide corresponding to the V3 domain (3) of MN-rgp120 was tested, it was found that none of the CD4 blocking antibodies showed any reactivity. However the epitope recognized by the non-CD4 blocking
35 monoclonal antibody, 1026, prepared against MN-rgp120 could be localized to the V3 domain by virtue of its

binding to this peptide. In other experiments, three synthetic peptides from the C4 domain of gp120 that incorporated sequences recognized by the CD4 blocking, weakly neutralizing monoclonal antibodies described by McKeating et al. (26) were tested (Table 4). It was found that none of the CD4 blocking monoclonal antibodies to MN-rgp120 reacted with these peptides, however the non-neutralizing, CD4 blocking 13H8 monoclonal antibody bound to the peptides corresponding to residues 423-440 of IIIB-gp120 and residues 431-441 of MN-gp120, but not to that corresponding to residues 426-437 of IIIB-gp120. Thus the 13H8 monoclonal antibody recognized an epitope that was similar, if not identical, to that described by McKeating et al. (26). This result is consistent with the observation that the 13H8 monoclonal antibody and the monoclonal antibodies described by Cordell et al. (4) and McKeating et al. (26) exhibited considerable cross reactivity, whereas the antibodies to MN-rgp120 were highly strain specific.

CD4 blocking antibodies recognize epitopes in the C4 domain. Previously, a strain specific, CD4 blocking monoclonal antibody (5C2) raised against IIIB-rgp120 was found to recognize an epitope in the C4 domain of IIIB-rgp120 (21, 33). Although the 5C2 monoclonal antibody was able to block the binding of rgp120 to CD4, it was unable to neutralize HIV-1 infectivity in vitro (7). Affinity columns prepared from 5C2 adsorbed an 11 amino acid peptide (residues 422 to 432) from a tryptic digest of gp120 (21), however monoclonal antibody 5C2 was unable to recognize this peptide coated onto wells of microtiter dishes in an ELISA format (Nakamura et al., unpublished results).

To determine whether the CD4 blocking monoclonal antibodies raised against MN-rgp120 recognized the

corresponding epitope in the C4 domain of MN-rgp120, a series of overlapping fragments, spanning the V4 and C4 domains of HIV-1_{MN} gp120, were prepared for expression in mammalian cells. A diagram of the fragments expressed is shown in Figures 3A and 3B. The C4 domain fragments were expressed as fusion proteins that incorporated the signal sequence and amino terminal 25 amino acids of HSV-1 glycoprotein D as described above.

Plasmids directing the expression of the chimeric C4 domain fragments were transfected into 293 cells, and their expression was monitored by radioimmunoprecipitation studies where a monoclonal antibody, 5B6, specific for the mature amino terminus of glycoprotein D was utilized. It was found (Figure 3B) that all of the fragments were expressed and exhibited mobilities on SDS-PAGE gels appropriate for their size. Thus fMN.368-408 (lane 1) exhibited a mobility of 19 kD; fMN.368-451 (lane 2) exhibited a mobility of 29 kD; fMN.419-433 (lane 3) exhibited a mobility of 6 kD, and fMN.414-451 (lane 4) exhibited a mobility of 6.1 kD.

The binding of monoclonal antibody 1024 to the recombinant fragments was then determined by ELISA (as described in Example 1). It was found (Figure 3A) that monoclonal antibody 1024 reacted with the fragments that contained the entire C4 domain of MN-rgp120 (fMN₃₆₈₋₄₅₁, fMN₄₀₄₋₄₅₅), but failed to bind to a fragment derived from the adjacent V4 domain (fMN₃₆₈₋₄₀₈) or to another fragment that contained V4 domain sequences and the amino terminal half of the C4 domain (fMN₃₆₈₋₄₂₈). The fact that 1024 bound to the fMN₄₁₄₋₄₅₁ and fMN₄₁₉₋₄₄₃ fragments demonstrated that the epitopes recognized by all of these monoclonal antibodies were contained entirely between residues 419 and 443 in the C4 domain.

35

Residues recognized by monoclonal antibodies that block binding of MN-rgp120 to CD4. To identify specific amino acid residues that might be part of the epitopes recognized by these monoclonal antibodies, the sequence of the C4 domain of MN-rgp120 was compared to those of the gp120s from the six other rgp120s that failed to react with the CD4 blocking monoclonal antibodies (Figure 4). It was noted that the sequence of MN-rgp120 was unique in that K occurred at position 429 whereas the other rgp120s possessed either E, G, or R at this position. Another difference was noted at position 440 where E replaced K or S. To evaluate the significance of these substitutions, a series of point mutations were introduced into the MN-rgp120 gene (Figure 5). Plasmids expressing the mutant proteins were transfected into 293s cells, and expression was verified by radioimmunoprecipitation with a monoclonal antibody (1034) directed to the V3 domain of MN-rgp120. Cell culture supernatants were harvested and used for the monoclonal antibody binding studies shown in Table 6. To verify expression, radioimmunoprecipitation studies using cell culture supernatants from cells metabolically labeled with [³⁵S]-methionine were performed using the 1024 monoclonal antibody specific for the C4 domain of MN-rgp120 (A) or the 1034 monoclonal antibody specific for the V3 domain of MN-rgp120. Immune complexes were precipitated with the use of fixed *S. aureus* and the adsorbed proteins were resolved by SDS-PAGE. Proteins were visualized by autoradiography. The samples were: Lane 1, MN.419A; lane 2 MN.421A; lane 3 MN.429E; lane 4, MN.429A; lane 5, MN.432A; lane 6, MN.440A; lane 7, MN-rgp120. The immunoprecipitation study showed that 1024 antibody binds well to all the variants except 3 and 4 which are mutated at residue 429. 1034 antibody

was used as a control and precipitates with anti-V3 antibodies.

The effect of these mutations on the binding of the CD4 blocking monoclonal antibodies was then evaluated by ELISA as illustrated in Table 5, below.

Table 5
Binding of CD4 blocking monoclonal antibodies to C4 domain mutants

Proteins/ MAbs	1024	1093	1096	1097	1110	1112	1127	5C2
MN-rgp120	1.0	1.0	1.0	1.0	1.0	1.0	1.0	0.05
MN-419A	1.11	1.10	0.94	1.21	0.78	0.95	1.10	ND
MN-421A	1.11	1.60	0.88	1.42	1.34	0.91	1.10	ND
MN-429E	0.03	0.07	0.11	0.04	0.10	0.10	0.02	ND
MN-429A	0.10	0.07	0.14	0.04	0.09	0.11	0.05	ND
MN-432A	0.77	0.15	0.59	0.08	0.12	0.24	0.26	ND
MN-440A	1.06	1.13	1.08	0.87	1.12	1.0	1.3	ND
IIIB-rgp120	0.03	ND	ND	ND	ND	ND	ND	1.0
MN-423F	ND	ND	ND	ND	ND	ND	ND	0.45
MN-423F, 429E	ND	ND	ND	ND	ND	ND	ND	1.09

Data represent the relative binding of MAbs to the native and mutant forms of rgp120. Values were calculated by dividing the binding (determined by ELISA) of the CD4 blocking MAbs to the proteins indicated by the values obtained for the binding of a V3 specific MAb (1034) to the same proteins (as described in Example 1).

It was found that replacement of K₄₄₀ with an A residue (MN.440A) had no effect on the binding of the 1024 monoclonal antibody or any of the other CD4 blocking monoclonal antibodies (Table 5). The significance of K at position 429 was then evaluated by substitution of either A (MN.429A) or E (MN.429E) at this location. It was found that the A for K substitution at position 429 (MN.420A) markedly reduced the binding of the 1024 monoclonal antibody and all of the other CD4 blocking monoclonal antibodies (Table 5).

Similarly, the replacement of E for K (MN.429E) at this position totally abrogated the binding of the 1024 monoclonal antibody and all of the other CD4 blocking monoclonal antibodies (Table 5). Several other mutants were constructed to evaluate the role of positively charged residues in the C4 domain. It was found that A for K substitutions at positions 419 (MN.419A) and 421 (MN.421A) failed to interfere with the binding of any of the CD4 blocking monoclonal antibodies as illustrated in Table 6, below.

Table 6
Correlation Between Antibody Binding Affinity
and Virus Neutralizing Activity

MAB	Block	K_d , nM ^c	IC_{50} , nM ^d
1024 ^e	+	2.7 ± 0.9	0.4
1086 ^{e,f}	-	9.7 ± 2.2	-
1093 ^e	+	9.9 ± 2.6	3.3
1096 ^e	+	10 ± 6	12
1097 ^e	+	13.4 ± 3.7	12
1110 ^e	+	12.1 ± 1.7	12
1112 ^e	+	20 ± 4.4	200
1127 ^e	+	9.3 ± 4	3.3
1086 ^{e,f}	-	9.7 ± 2.2	-
13H8 ^{f,g}	+ ^b	22 ± 6	-

^a Blocked binding of rgp120 MN to CD4.

^b Blocked binding of rgp120 IIIb, not rgp120 MN, to CD4.

^c Mean of four determinations calculated using the method of Scatchard (40).

^d Neutralization of HIV-1_{MN} infectivity in vitro.

^e Anti-rgp120 MN antibody.

^f Did not neutralize HIV-1 infectivity.

^g Anti-rgp120 IIIb antibody.

However, when K at position 432 was replaced with A (MN432.A), the binding of all of the CD4 blocking antibodies was markedly reduced (Table 5). Interestingly, the binding of monoclonal antibody 1024 appeared less affected by this substitution than the other monoclonal antibodies (Table 5). Thus, these studies demonstrated that K₄₂₉ and K₄₃₂ were critical for the binding of all of the CD4 blocking monoclonal antibodies, and that K₄₁₉, K₄₂₁, and K₄₄₀ did not appear to play a role in monoclonal antibody binding.

Amino acids recognized monoclonal antibodies that block binding of IIIB-rgp120 to CD4. The identification of residues 429 and 432 as being part of the epitope recognized by the MN-rgp120 specific CD4 blocking monoclonal antibodies was particularly interesting since this region was previously found to be implicated in the binding of the 5C2 monoclonal antibody (21). The properties of the 1024 like-monoclonal antibodies and the 5C2 monoclonal antibody differed from the C4 reactive monoclonal antibodies described by other investigators (4, 43) in that the former appeared strain specific and the latter were broadly cross reactive. To account for the strain specificity of these monoclonal antibodies, the sequence of the eleven amino acid peptide of IIIB-rgp120 recognized by monoclonal antibody 5C2 was compared to the corresponding sequence of MN-rgp120. It was found that the IIIB protein differed from the MNB protein at positions 429 where K replaced E and at position 423 where I replaced F (Figure 5). Because it was known from previous studies (33) that the 5C2 monoclonal antibody was unable to bind to gp120 from two strains (i.e., NY-5 and JRcsf) that also possessed E at position 423, it seemed unlikely that this position could account for the strain specificity of

5C2. Sequence comparison (Figure 5) also showed that gp120 from HIV-1_{mb} was unique in that a phenylalanine residue occurred at position 423 whereas the other six strains examined possess an I at this position.

5 To determine whether residues 423 and/or 429 could account for the type specificity of the 5C2 monoclonal antibody, a mutant of MN-rgp120 was constructed which incorporated an F for I replacement at position 423 (MN.423F). In addition, the MN-rgp120 mutant, MN.429E
10 (described above) was further mutagenized to incorporate a F for I substitution at position 423 (MN.423F), thus resulting in a double mutant (MN.423F,429E) whose sequence was identical to that of IIIB-rgp120 within the 10 amino acid 5C2 epitope
15 (Figure 4). The expression of these mutants in 293s cells was verified by radioimmunoprecipitation using rabbit polyclonal antisera to MN-rgp120. When the binding of the 13H8 monoclonal antibody to a set of mutants incorporating substitutions at position 423 and
20 429 was examined, it was found that none of the replacements effected the binding of this antibody (data not shown). When the 5C2 monoclonal antibody was examined, it was found that the F for I replacement (MN.423 F) conferred partial reactivity (Table 5).
25 When the double mutant (MN.423F,429E), containing the F for I substitution as well as the E for K substitution was tested, binding that was indistinguishable from that to IIIB-rgp120 was observed (Table 5). These results demonstrated that F at position 423 and E at
30 position 429 both play a role in binding of the 5C2 monoclonal antibody, and suggest that the strain specificity of 5C2 can be attributed to the residues at these positions.

Examination of the sequences of gp120 from the
35 various clones of LAI that have been analyzed revealed that several substrains of LAI differed from each other

in the C4 domain. Thus the sequences of the IIIB (30), Bru (46), and HXB3 (6) clones of LAI were identical at positions 423 and 429 where F and E residues occurred respectively. However, the sequence of the HXB2

5 substrain (36) differed from the others at these positions where, like MN-rgp120, K replaced E and at position 423 where I replaced F (Figure 5). Similarly, the HX10 and BH10 substrains (36, 37) differed only at position 423 where, like HIV-1_{MN}, I replaced F. Based

10 on the mutagenesis experiments above, it would be predicted that monoclonal antibody 1024 should be able to bind to gp120 from the HXB2 substrain of LAI, but not the HXB3 substrain. If I₄₂₃ was important for binding, then 1024 should also bind the HX10 substrain.

15 To test this hypothesis, the binding of monoclonal antibody 1024 to the surface cells infected with either IIIB, HXB2, HXB3, and HX10 substrains of HIV-1_{LAI} was measured by flow cytometry. It was found that monoclonal antibody 1024 was able to bind only HXB2

20 providing further confirmation that residues 423 and 429 were important for the binding of this antibody. The fact that monoclonal antibody 1024 did not bind to HX10 infected cells suggested that I₄₂₃ was not important for the binding of this monoclonal antibody. Thus

25 these studies demonstrate that reactivity with the 1024 monoclonal antibody segregates with the occurrence of F and E residues at positions 423 and 429, respectively, and shows that substrains of HIV-1_{LAI} differ from one another at a functionally significant epitope in the C4

30 domain.

Neutralizing activity of CD4 blocking antibodies correlates with their binding affinity. To account for the difference in virus neutralizing activity between

35 the CD4 blocking monoclonal antibodies, their gp120 binding affinities were determined by competitive

binding of [125 I]-labeled monoclonal antibody to rgp120 (Table 6). Typical Scatchard analysis of data from these assays is shown in Figure 7 (A to C). Linear, one-site binding kinetics were observed for all the monoclonal antibodies to MN-rgp120, suggesting that only a single class of sites was recognized, and that there was no cooperativity between two combining sites of each immunoglobulin molecule. It was found (Figure 7A, Table 6) that monoclonal antibody 1024, which exhibited the most potent virus neutralizing activity (IC_{50} of 0.08 μ g per ml), possessed the lowest K_d (2.7 nM). In contrast (Figure 7C, Table 6), monoclonal antibody 1112, the antibody that exhibited the weakest virus neutralizing activity (IC_{50} of 30 μ g per ml) possessed the highest K_d (20 nM). K_d s for six additional CD4-blocking monoclonal antibodies raised against MN-rgp120 were also determined (Table 6). It was found that monoclonal antibodies that possessed intermediate K_d s similarly possessed intermediate neutralization IC_{50} values. To explore the relationship between virus neutralizing activity and gp120 binding affinity, the data in Table 6 was plotted in several different ways. It was found that when the K_d of the monoclonal antibodies was plotted as a function of the log of the IC_{50} , a linear relationship was obtained (Figure 8). Using this analysis a correlation coefficient (r) of 0.97) was obtained. Thus, this graph demonstrates that the virus neutralizing activity of these monoclonal antibodies is directly proportional to the gp120 binding affinity, and that the threshold for neutralization at this epitope is defined by the slope of the graph in Figure 8.

A similar analysis was performed with the non-neutralizing CD4 blocking monoclonal antibodies to IIIIB-rgp120, 5C2 and 13H8. The binding curve for 13H8 (Figure 7C) showed that it bound to a single class of

sites on IIIB-rgp120 with a K_d of 22 nM. The affinity of 5C2 could not be determined by this assay because at antibody concentrations greater than 5 nM, non-linear (reduced gp120 binding) was observed. This effect was suggestive steric hindrance at these concentrations or negative cooperativity between combining sites. The binding affinity was also determined for the non-neutralizing, non-CD4 blocking monoclonal antibody to MN-rgp120, 1086. The fact that this antibody exhibited a binding affinity similar (9.7 nM) to many of the neutralizing monoclonal antibodies but failed to inhibit infectivity, proves that high antibody binding affinity alone is not sufficient for neutralization.

Effect of C4 Domain Mutants on CD4 binding.

Finally, the CD4 binding properties of the series of MN-rgp120 mutants, constructed to localize the C4 domain epitopes, were measured in a qualitative co-immunoprecipitation assay. In these studies the ability of the mutagenized MN-rgp120 variants to co-immunoprecipitate CD4 was evaluated as described previously (21) in a qualitative co-immunoprecipitation assay similar to that described previously (19). Briefly, 293 cells, transfected with plasmids directing the expression of MN-rgp120 variants described in Figure 5, were metabolically labeled with [35 S]-methionine, and the growth conditioned cell culture supernatants were incubated with rsCD4. The resulting rsCD4:gp120 complexes were then immunoprecipitated by addition of the CD4 specific monoclonal antibody, 465 (A) or a positive control monoclonal antibody (1034) directed to the V3 domain of MN-rgp120 (B). The immunoprecipitated proteins were resolved by SDS-PAGE and visualized by autoradiography as described previously (3). The samples were: Lane 1, MN.419A; lane 2, MN.421A; lane 3, MN.429E; lane 4,

MN.429A; lane 5, MN.432A; lane 6, MN.440A; lane 7, MN-rgp120. The gel showed that the mutants that block antibody binding do not block binding of CD4. Therefore, the antibodies do not bind to the gp120 CD4-binding contact residues. This indicates that steric hinderance may inhibit antibody binding, rather than that the antibodies bind directly to the CD4 contact residues to inhibit binding.

It was found that all of the variants in which apolar A residue was substituted for the charged K or E residues (e.g., MN.419A, MN.421A, MN.432A, and MN.440A) were still able to co-immunoprecipitate rsCD4. Similarly, the replacement of E for K at position 429 (MN.429E), the replacement of F for I at position 423 (MN.423F) or the mutant which incorporated both mutation (MN.423F,429E) also showed no reduction in their ability to co-immunoprecipitate rsCD4. Thus, radical amino acid substitutions at five positions failed to affect the binding of gp120 to CD4. These results were consistent with previous studies (5, 21, 34) where it was found that only a few of the many mutations that have been induced in this region effected CD4 binding.

This study indicates that neutralizing epitopes in the C4 domain have now been found to be located between about residues 420 and 440. In addition, the critical residues for antibody binding are residues 429 and 432.

EXAMPLE 2

Identification of V2 Neutralizing Epitopes

The procedures described in Example 1 were used to map epitopes in the V2 region of gp120. Table 7 illustrates the results of mutagenicity studies to map V2 neutralizing epitopes. In the table, the columns indicate the comparison of binding of the monoclonal antibodies with wild type (WT) gp120 in comparison to

various mutations of gp120 using standard notation.

For example, "G171R" indicates that the glycine (G) at residue 171 has been replaced by an arginine (R).

"172A/173A" indicates that the residues at 172 and 173

5 have been replaced by alanine. The neutralizing monoclonal antibodies tested (MAbs) are listed in the rows. The numerical values in the table are the optical density value of an ELISA assay performed as described in Example 1 to measure the amount of
10 antibody binding. The underlined values indicate significantly reduced binding, indicating the substituted residue is critical for binding of the antibody.

15

TABLE 7

	WT	G171R, M174V	172A/ 173A	E187V	187V/188S
<u>MAbs</u>					
6E10	1.00	<u>0.10</u>	1.28	0.60	<u>0.25</u>
1017	1.00	0.70	1.10	0.87	<u>0.04</u>
20 1022	1.00	0.80	1.10	1.00	<u>0.00</u>
1028	1.00	0.90	1.18	1.07	<u>0.04</u>
1029	1.00	0.83	1.16	1.01	<u>0.16</u>
1019	1.00	<u>0.13</u>	1.30	0.75	0.74
1027	1.00	<u>0.00</u>	1.20	0.80	0.64
25 1025	1.00	0.69	<u>0.00</u>	<u>0.00</u>	0.83
1088	1.00	0.73	1.12	0.94	<u>0.03</u>
13H8	1.00	0.77	0.78	0.48	0.65

TABLE 7 (continued)

	WT	177A	172A/173A	188A	183A
<u>MAbs</u>					
6E10	1.00	<u>0.36</u>	0.52	0.64	0.43
5 1017	1.00	0.77	0.77	0.76	<u>0.11</u>
1022	1.00	0.86	0.72	<u>0.14</u>	<u>0.00</u>
1028	1.00	0.93	0.78	0.49	<u>0.04</u>
1029	1.00	0.88	0.85	0.53	<u>0.16</u>
1019	1.00	<u>0.16</u>	<u>0.00</u>	0.41	0.44
10 1027	1.00	<u>0.00</u>	<u>0.02</u>	0.41	0.49
1025	1.00	0.75	0.0	0.83	0.72
1088	1.00	0.77	0.77	0.53	<u>0.00</u>
13H8	1.00	0.72	0.72	0.53	0.60

15 As illustrated in Table 7, the study demonstrated that there are a series of overlapping neutralizing epitopes from been found to be located in the V2 region (residues 163 through 200), with most of the epitopes located between residues 163 and 200. In addition, the study indicates that the critical residues in the V2 domain for antibody binding are residues 171, 173, 174, 177, 181, 183, 187, and 188.

EXAMPLE 3

25 Immunization Studies

gp120 from the MN, GNE₃, and GNE₁₆ strains of HIV was prepared by amplifying the gene from each isolate and cloning and expressing the gene in CHO cells as described in Berman et al., *J. Virol.* 66:4464-4469 (1992). Briefly, the gp160 gene was amplified with two rounds of amplification using the following nested

primers according to the protocol by Kellog et al., pp 337-347 in *PCR Protocols: a guide to methods and amplification*. Innis et al. (eds.) Academic Press, Inc., New York.

5 First round primers:

AATAATAGCAATAGTTGTGTGGWCC (W is A or T)

ATTCTTTCCCTTAYAGTAGGCCATCC (Y is T or C)

Second round primers:

GGGAATTCGGATCCAGAGCAGAAGACAGTGGCAATGA

10 GTCAAGAATTCTTATAGCAAAGCCCTTTCCAA

The primers are SEQ. ID. NOS. 31-34. Each gene is then digested with the restriction endonucleases KpnI and AccI. The resulting fragment was subcloned into the Bluescript (+) phagemid M13 vector (Stratagene, Inc.) and sequenced by the dideoxynucleotide method (Sanger et al., *Proc. Natl. Acad. Sci. USA* 74:5463-5467 (1977)).

A fragment of the gp120 coding region was then used to construct a chimeric gene for expression in mammalian cells, as described in Lasky et al., *Science* 223:209-212 (1986). The 5' end was fused to a polylinker adjacent to a simian virus 40 (SV40) promoter and the 3' end was fused to a polylinker adjacent to the 3' untranslated sequences containing an SV40 polyadenylation signal. The expression vector (MN-rgp120) was co-transfected in CHO cells deficient in production of the enzyme dihydrofolate reductase, along with a plasmid (pSVdhfr) containing a cDNA encoding the selectable marker, dihydrofolate reductase. Cell lines expressing MN-rgp120 were isolated as described in Lasky et al., *Science* 223:209-212 (1986). The recombinant glycoprotein was purified from growth-conditioned cell culture medium by immunoaffinity and ion exchange chromatography as described in Leonard et al., *J. Biol. Chem.* 265:10373-10382 (1990).

gp120 from the GNE₈ and GNE₁₆ strains of HIV is prepared in the same manner as described for the MN isolate.

5 MN-rgp120 (300 µg/injection), GNE₈-rgp120 (300 µg/injection), and GNE₁₆-rgp120 (300 µg/injection) are prepared in an aluminum hydroxide adjuvant (as described in Cordonnier et al., *Nature* 340:571-574 (1989)). Six chimpanzees are injected at 0, 4, and 32 weeks. Sera are collected and assayed for neutralizing
10 antibody to each strain of HIV at the time of each immunization and three weeks thereafter. At 35 weeks, each of the chimpanzees has significant levels of neutralizing antibodies to each strain.

At 35 weeks, the chimpanzees are randomly assigned
15 to three groups. Each group is challenged with about 10 50% chimpanzee-infectious doses (CID₅₀) each of one of the vaccine isolates. One unimmunized chimpanzee (control) is also injected with the same amount of virus as the immunized chimpanzees for each vaccine
20 strain.

Sera are drawn every two weeks throughout the study and assayed for antibodies to HIV core proteins and for the presence of HIV by PCR amplification and co-cultivation of peripheral blood mononuclear cells
25 (PBMCs) from the chimpanzee together with activated human or chimpanzee PBMCs. The presence of antibodies to core proteins indicates the presence of viral infection as does the detection of amplified viral DNA or viral infection of co-cultivated cells.

30 The presence of virus is detected by PCR and co-cultivation methods in each unimmunized control animal between weeks 2 and 4 post challenge. Antibodies to core proteins appear in the control chimpanzees at six weeks post challenge. Neither virus
35 nor antibodies are at detectable levels in any of the immunized chimpanzees at one year post challenge,

indicating that the vaccine effectively protects the chimpanzees from infection from each of the challenge strains.

REFERENCES

1. Berman, P.W. et al., 1989. Expression and immunogenicity of the extracellular domain of the human immunodeficiency virus type 1 envelope glycoprotein, gp160. J. Virol. 63:3489-3498.
2. Berman, P. W. et al., 1990. Protection of chimpanzees from infection by HIV-1 after vaccination with gp120 but not gp160. Nature 345:622-625.
3. Berman, P. W. et al., 1992. Neutralization of multiple laboratory and clinical isolates of HIV-1 by antisera raised against gp120 from the MN isolate of HIV-1. J. Virol. 7:4464-4469.
4. Cordell, J. et al., 1991. Rat monoclonal antibodies to non-overlapping epitopes of human immunodeficiency virus type 1 gp120 block CD4 binding in vitro. Virology 185:72-79.
5. Cordonnier, A. et al., 1989. Single amino acid changes in HIV envelope affect viral tropism and receptor binding. Nature 340:571-574.
6. Crowl, R. et al., 1985. HTLV-III env gene products synthesized in E. coli are recognized by antibodies present in the sera of AIDS patients. Cell 41:979-986.
7. Dowbenko, D. et al., 1988. Epitope mapping of the human immunodeficiency virus type 1 gp120 with monoclonal antibodies. J. Virol. 62:4703-4711.
8. Eaton, D. et al., 1986. Construction and characterization of an active factor VIII lacking the central one-third of the molecule. Biochemistry 29:8343-8347.
9. Fouchier, R. A. M. et al., 1992. Phenotype-associated sequence variation in the third variable domain of the human immunodeficiency virus type 1 gp120 molecule. J. Virol. 66: 3183-3187.
10. Goudsmit, J. et al., 1988. Human immunodeficiency virus type 1 neutralization epitope with conserved architecture elicits early type-specific antibodies in

- experimentally infected chimpanzees. *Proc. Natl. Acad. Sci. U.S.A.* 85:4478-4482.
11. Graham, F. et al., 1973. A new technique for the assay of infectivity of human adenovirus 5 DNA. *Virology* 52:456-467.
12. Graham, F. L. et al., 1977. Characteristics of a human cell line transformed by the human adenovirus type 5. *J. Gen. Virol.* 36:59-77.
13. Gurgo, C. et al., 1988. Envelope sequences of two new United States HIV-1 isolates. *Virology* 164: 531-536.
14. Haffar, O.K. et al., 1991. The cytoplasmic tail of HIV-1 gp160 contains regions that associate with cellular membranes. *Virology* 180:439-441.
15. Higuchi, R. 1990. Recombinant PCR. p.177-183. In M. A. Innis et al. (eds.), *PCR Protocols A Guide to Methods and Applications*, Academic Press, Inc., New York.
16. Ho, D. D. et al., 1991. Conformational epitope on gp120 important in CD4 binding and human immunodeficiency virus type 1 neutralization identified by a human monoclonal antibody. *J. Virol.* 65:489-493.
17. Kellog, D. E. et al., 1990. Detection of Human Immunodeficiency Virus, p. 337-347. In M. A. Innis et al. (eds.), *PCR Protocols A Guide to Methods and Applications*, Academic Press, Inc., New York.
18. Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680-685.
19. Langedijk, J. P. M. et al., 1991. Neutralizing activity of anti-peptide antibodies against the principal neutralization domain of human immunodeficiency virus type 1. *J. Gen. Virol.* 72:2519-2526.
20. LaRosa, G. J. et al., 1990. Conserved sequences and structural elements in the HIV-1 principal neutralizing determinant. *Science* 249:932-935.

21. Lasky, L. A. et al., 1987. Delineation of a region of the human immunodeficiency virus gp120 glycoprotein critical for interaction with the CD4 receptor. *Cell* 50:975-985.
- 5 22. Lasky, L. A. et al., 1986. Neutralization of the AIDS retrovirus by antibodies to a recombinant envelope glycoprotein. *Science* 233: 209-212.
23. Matsushita, S. et al., 1988. Characterization of a human immunodeficiency virus neutralizing monoclonal antibody and mapping of a neutralizing epitope. *J. Virol.* 62:2107-2114.
- 10 24. McCutchan, F.E. et al., 1992. Genetic Variants of HIV-1 in Thailand. *AIDS Res. and Human Retroviruses* 8:1887-1895.
- 15 25. McKeating, J. et al., 1991. Recombinant CD4-selected human immunodeficiency virus type 1 variants with reduced gp120 affinity for CD4 and increased cell fusion capacity. *J. Virol.* 65: 4777-4785.
26. McKeating, J. A. et al., 1992. Monoclonal antibodies to the C4 region of human immunodeficiency virus type 1 gp120: use in topological analysis of a CD4 binding site. *AIDS Research and Human Retroviruses*. 8: 451-459.
- 20 27. McNearney, T. et al., 1992. Relationship of human immunodeficiency virus type 1 sequence heterogeneity to stage of disease. *Proc. Natl. Acad. Sci. U.S.A.* 89:10247-10251.
28. Modrow, S. et al., 1987. Computer-assisted analysis of envelope protein sequences of seven human immunodeficiency virus isolates: predictions of antigenic epitopes in conserved and variable regions. *J. Virol.* 61:570-578.
- 30 29. Moore, J. P. 1990. Simple methods for monitoring HIV-1 and HIV-2 gp120 binding to sCD4 by ELISA: HIV-1 has a 25 fold lower affinity than HIV-1 for sCD4. *AIDS* 3:297-305.
- 35

30. Muesing, M. A. et al., 1985. Nucleic acid structure and expression of the human AIDS/lymphadenopathy retrovirus. *Nature* 313:450-458.
31. Munson, P.J. et al. 1983. LIGAND: a computerized analysis of ligand binding data. *Methods Enzymol.* 92:543.
32. Myers, G. et al., 1992. Human Retroviruses and AIDS. A compilation and analysis of nucleic acid and amino acid sequences. Los Alamos National Laboratory, Los Alamos, New Mexico.
33. Nakamura, G. et al., 1992. Monoclonal antibodies to the extracellular domain of HIV-1_{MB} gp160 that neutralize infectivity, block binding to CD4, and react with diverse isolates. *AIDS and Human Retroviruses* 8:1875-1885.
34. Olshevsky V. et al., 1990. Identification of individual human immunodeficiency virus type 1 gp120 amino acids important for CD4 receptor binding. *J. Virol.* 64:5701-5707.
35. Posner, M. R. et al., 1991. An IgG human monoclonal antibody which reacts with HIV-1/GP120, inhibits virus binding to cells and neutralizes infection. *J. Immunol.* 146:4325-4332.
36. Ratner, L. et al., 1987. Complete nucleotide sequences of functional clones of the AIDS virus. *AIDS Res. and Human Retroviruses* 3:57-69.
37. Ratner, L. et al., 1985. Complete nucleotide sequence of the AIDS virus, HTLV-III. *Nature* 313:277-284.
38. Reitz, M.S. Jr. et al., 1992. On the historical origins of HIV-1 (MN) and (RF). *AIDS Research and Human Retroviruses* 9: 1539-1541.
39. Rusche, J.R. et al., 1988. Antibodies that inhibit fusion of human immunodeficiency virus-infected cells bind to a 24-amino acid sequence of the viral envelope, gp120. *Proc. Natl. Acad. Sci. USA.* 85:3198-3202.

40. Scatchard, G. 1949. The attractive proteins for small molecules and ions. Ann. N.Y. Acad. Sci. 51: 660-672.
41. Schnittman, S. M. et al., 1988. Characterization of gp120 binding to CD4 and an assay that measures ability of sera to inhibit this binding. J. Immunol. 141:4181-4186.
42. Scott, C.F. Jr. et al., 1990. Human monoclonal antibody that recognizes the V3 region of human immunodeficiency virus gp120 and neutralizes the human T-lymphotropic virus type III_{MN} strain. Proc. Natl. Acad. Sci. U.S.A. 87:8597-8601.
43. Sun, N. C. et al., 1989. Generation and characterization of monoclonal antibodies to the putative CD4-binding domain of human immunodeficiency virus type 1 gp120. J. Virol. 63:3579-3585.
44. Tersmette, M. R. A. et al., 1989. Evidence for a role of virulent human immunodeficiency virus (HIV) variants in the pathogenesis of AIDS obtained from studies on a panel of sequential HIV isolates. J. Virol. 63: 2118-2125.
45. Tilley, S. A. et al., 1991. A human monoclonal-antibody against the CD4-binding site of HIV-1 GP120 exhibits potent, broadly neutralizing activity. Res. Virology 142:247-259.
46. Wain Hobson, S. et al., 1985. Nucleotide sequence of the AIDS virus, LAV. Cell 40:9-17.
47. Weiss, R. A. et al., 1986. Variable and conserved neutralizing antigen of human immunodeficiency virus. Nature 324:572-575.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT: Berman, Phillip W.
Nakamura, Gerald R.

(ii) TITLE OF INVENTION: HIV Envelope Polypeptides

10

(iii) NUMBER OF SEQUENCES: 26

(iv) CORRESPONDENCE ADDRESS:

15

(A) ADDRESSEE: Skjerven, Morrill, MacPherson, Franklin &
Friel

(B) STREET: 25 Metro Drive Suite 700

(C) CITY: San Jose

(D) STATE: California

(E) COUNTRY: U.S.A.

20

(F) ZIP: 95110

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

25

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

30

(B) FILING DATE:

(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Terlizzi, Laura

35

(B) REGISTRATION NUMBER: 31,307

(C) REFERENCE/DOCKET NUMBER: M-2820-1P

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (408) 283-1222

40

(B) TELEFAX: (408) 283-1233

(2) INFORMATION FOR SEQ ID NO:1:

45

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 511 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

	Met	Arg	Val	Lys	Gly	Ile	Arg	Arg	Asn	Tyr	Gln	His	Trp	Trp	Gly	Arg	
	1				5					10					15		
5	Gly	Thr	Met	Leu	Leu	Gly	Leu	Leu	Met	Ile	Cys	Ser	Ala	Thr	Glu	Lys	
				20					25					30			
	Leu	Trp	Val	Thr	Val	Tyr	Tyr	Gly	Val	Pro	Val	Trp	Lys	Glu	Ala	Thr	
			35					40					45				
10	Thr	Thr	Leu	Phe	Cys	Ala	Ser	Asp	Ala	Lys	Ala	Tyr	Asp	Thr	Glu	Ala	
			50				55						60				
	His	Asn	Val	Trp	Ala	Thr	His	Ala	Cys	Val	Pro	Thr	Asp	Pro	Asn	Pro	
	65					70					75				80		
15	Gln	Glu	Val	Glu	Leu	Val	Asn	Val	Thr	Glu	Asn	Phe	Asn	Met	Trp	Lys	
				85						90					95		
	Asn	Asn	Met	Val	Glu	Gln	Met	His	Glu	Asp	Ile	Ile	Ser	Leu	Trp	Asp	
20				100					105					110			
	Gln	Ser	Leu	Lys	Pro	Cys	Val	Lys	Leu	Thr	Pro	Leu	Cys	Val	Thr	Leu	
			115					120					125				
25	Asn	Cys	Thr	Asp	Leu	Arg	Asn	Thr	Thr	Asn	Thr	Asn	Asn	Ser	Thr	Asp	
		130					135					140					
	Asn	Asn	Asn	Ser	Lys	Ser	Glu	Gly	Thr	Ile	Lys	Gly	Gly	Glu	Met	Lys	
	145				150						155				160		
30	Asn	Cys	Ser	Phe	Asn	Ile	Thr	Thr	Ser	Ile	Gly	Asp	Lys	Met	Gln	Lys	
				165					170						175		
	Glu	Tyr	Ala	Leu	Leu	Tyr	Lys	Leu	Asp	Ile	Glu	Pro	Ile	Asp	Asn	Asp	
35				180					185					190			
	Ser	Thr	Ser	Tyr	Arg	Leu	Ile	Ser	Cys	Asn	Thr	Ser	Val	Ile	Thr	Gln	
			195					200					205				
40	Ala	Cys	Pro	Lys	Ile	Ser	Phe	Glu	Pro	Ile	Pro	Ile	His	Tyr	Cys	Ala	
		210					215					220					
	Pro	Ala	Gly	Phe	Ala	Ile	Leu	Lys	Cys	Asn	Asp	Lys	Lys	Phe	Ser	Gly	
	225				230						235				240		
45	Lys	Gly	Ser	Cys	Lys	Asn	Val	Ser	Thr	Val	Gln	Cys	Thr	His	Gly	Ile	
				245					250					255			
	Arg	Pro	Val	Val	Ser	Thr	Gln	Leu	Leu	Leu	Asn	Gly	Ser	Leu	Ala	Glu	
50				260				265						270			
	Glu	Glu	Val	Val	Ile	Arg	Ser	Glu	Asp	Phe	Thr	Asp	Asn	Ala	Lys	Thr	

	275	280	285
	Ile Ile Val His Leu Lys Glu Ser Val Gln Ile Asn Cys Thr Arg Pro		
	290	295	300
5	Asn Tyr Asn Lys Arg Lys Arg Ile His Ile Gly Pro Gly Arg Ala Phe		
	305	310	315
	Tyr Thr Thr Lys Asn Ile Lys Gly Thr Ile Arg Gln Ala His Cys Ile		
10		325	330
	Ile Ser Arg Ala Lys Trp Asn Asp Thr Leu Arg Gln Ile Val Ser Lys		
		340	345
			350
15	Leu Lys Glu Gln Phe Lys Asn Lys Thr Ile Val Phe Asn Pro Ser Ser		
		355	360
	Gly Gly Asp Pro Glu Ile Val Met His Ser Phe Asn Cys Gly Gly Glu		
		370	375
20			380
	Phe Phe Tyr Cys Asn Thr Ser Pro Leu Phe Asn Ser Ile Trp Asn Gly		
		385	390
			395
	Asn Asn Thr Trp Asn Asn Thr Thr Gly Ser Asn Asn Asn Ile Thr Leu		
25		405	410
			415
	Gln Cys Lys Ile Lys Gln Ile Ile Asn Met Trp Gln Lys Val Gly Lys		
		420	425
			430
30	Ala Met Tyr Ala Pro Pro Ile Glu Gly Gln Ile Arg Cys Ser Ser Asn		
		435	440
			445
	Ile Thr Gly Leu Leu Leu Thr Arg Asp Gly Gly Glu Asp Thr Asp Thr		
		450	455
			460
35	Asn Asp Thr Glu Ile Phe Arg Pro Gly Gly Gly Asp Met Arg Asp Asn		
		465	470
			475
	Trp Arg Ser Glu Leu Tyr Lys Tyr Lys Val Val Thr Ile Glu Pro Leu		
40		485	490
			495
	Gly Val Ala Pro Thr Lys Ala Lys Arg Arg Val Val Gln Arg Glu		
		500	505
			510

(25) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 501 amino acids

(B) TYPE: amino acid

50 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

	Lys	Tyr	Ala	Leu	Ala	Asp	Ala	Ser	Leu	Lys	Met	Ala	Asp	Pro	Asn	Arg	
	1				5					10					15		
5	Ph	Arg	Gly	Lys	Asp	Leu	Pro	Val	Leu	Asp	Gln	Leu	Leu	Glu	Val	Pro	
				20					25					30			
	Val	Trp	Lys	Glu	Ala	Thr	Thr	Thr	Leu	Phe	Cys	Ala	Ser	Asp	Ala	Lys	
10			35					40					45				
	Ala	Tyr	Asp	Thr	Glu	Ala	His	Asn	Val	Trp	Ala	Thr	His	Ala	Cys	Val	
	50						55					60					
15	Pro	Thr	Asp	Pro	Asn	Pro	Gln	Glu	Val	Glu	Leu	Val	Asn	Val	Thr	Glu	
	65					70				75						80	
	Asn	Phe	Asn	Met	Trp	Lys	Asn	Asn	Met	Val	Glu	Gln	Met	His	Glu	Asp	
				85					90						95		
20	Ile	Ile	Ser	Leu	Trp	Asp	Gln	Ser	Leu	Lys	Pro	Cys	Val	Lys	Leu	Thr	
				100					105					110			
	Pro	Leu	Cys	Val	Thr	Leu	Asn	Cys	Thr	Asp	Leu	Arg	Asn	Thr	Thr	Asn	
25			115					120					125				
	Thr	Asn	Asn	Ser	Thr	Asp	Asn	Asn	Asn	Ser	Lys	Ser	Glu	Gly	Thr	Ile	
	130						135						140				
30	Lys	Gly	Gly	Glu	Met	Lys	Asn	Cys	Ser	Phe	Asn	Ile	Thr	Thr	Ser	Ile	
	145					150					155					160	
	Gly	Asp	Lys	Met	Gln	Lys	Glu	Tyr	Ala	Leu	Leu	Tyr	Lys	Leu	Asp	Ile	
				165						170					175		
35	Glu	Pro	Ile	Asp	Asn	Asp	Ser	Thr	Ser	Tyr	Arg	Leu	Ile	Ser	Cys	Asn	
				180					185					190			
	Thr	Ser	Val	Ile	Thr	Gln	Ala	Cys	Pro	Lys	Ile	Ser	Phe	Glu	Pro	Ile	
40			195					200					205				
	Pro	Ile	His	Tyr	Cys	Ala	Pro	Ala	Gly	Phe	Ala	Ile	Leu	Lys	Cys	Asn	
	210						215					220					
45	Asp	Lys	Lys	Phe	Ser	Gly	Lys	Gly	Ser	Cys	Lys	Asn	Val	Ser	Thr	Val	
	225					230					235					240	
	Gln	Cys	Thr	His	Gly	Ile	Arg	Pro	Val	Val	Ser	Thr	Gln	Leu	Leu	Leu	
				245						250				255			
50	Asn	Gly	Ser	Leu	Ala	Glu	Glu	Glu	Val	Val	Ile	Arg	Ser	Glu	Asp	Phe	
				260					265					270			

Thr Asp Asn Ala Lys Thr Ile Ile Val His Leu Lys Glu Ser Val Gln
 275 280 285
 5 Ile Asn Cys Thr Arg Pro Asn Tyr Asn Lys Arg Lys Arg Ile His Ile
 290 295 300
 Gly Pro Gly Arg Ala Phe Tyr Thr Thr Lys Asn Ile Lys Gly Thr Ile
 305 310 315 320
 10 Arg Gln Ala His Cys Ile Ile Ser Arg Ala Lys Trp Asn Asp Thr Leu
 325 330 335
 Arg Gln Ile Val Ser Lys Leu Lys Glu Gln Phe Lys Asn Lys Thr Ile
 340 345 350
 15 Val Phe Asn Pro Ser Ser Gly Gly Asp Pro Glu Ile Val Met His Ser
 355 360 365
 20 Phe Asn Cys Gly Gly Glu Phe Phe Tyr Cys Asn Thr Ser Pro Leu Phe
 370 375 380
 Asn Ser Ile Trp Asn Gly Asn Asn Thr Trp Asn Asn Thr Thr Gly Ser
 385 390 395 400
 25 Asn Asn Asn Ile Thr Leu Gln Cys Lys Ile Lys Gln Ile Ile Asn Met
 405 410 415
 Trp Gln Lys Val Gly Lys Ala Met Tyr Ala Pro Pro Ile Glu Gly Gln
 420 425 430
 30 Ile Arg Cys Ser Ser Asn Ile Thr Gly Leu Leu Leu Thr Arg Asp Gly
 435 440 445
 Gly Glu Asp Thr Asp Thr Asn Asp Thr Glu Ile Phe Arg Pro Gly Gly
 450 455 460
 35 Gly Asp Met Arg Asp Asn Trp Arg Ser Glu Leu Tyr Lys Tyr Lys Val
 465 470 475 480
 40 Val Thr Ile Glu Pro Leu Gly Val Ala Pro Thr Lys Ala Lys Arg Arg
 485 490 495
 Val Val Gln Arg Glu
 500

45
 (2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- 50 (A) LENGTH: 28 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

5 Cys Lys Ile Lys Gln Ile Ile Asn Met Trp Gln Lys Val Gly Lys Ala
1 5 10 15
Met Tyr Ala Pro Pro Ile Glu Gly Gln Ile Arg Cys
20 25

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 28 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

20 Cys Arg Ile Lys Gln Phe Ile Asn Met Trp Gln Glu Val Gly Lys Ala
1 5 10 15
Met Tyr Ala Pro Pro Ile Ser Gly Gln Ile Arg Cys
25 20 25

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- 30 (A) LENGTH: 28 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

35 Cys Arg Ile Lys Gln Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala
1 5 10 15
40 Met Tyr Ala Pro Pro Ile Lys Gly Gln Ile Arg Cys
20 25

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 28 amino acids
(B) TYPE: amino acid
50 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

1 Cys Arg Ile Lys Gln Ile Ile Asn Met Trp Gln Gly Val Gly Lys Ala
5 1 5 10 15
20 Met Tyr Ala Pro Pro Ile Glu Gly Gln Ile Asn Cys
25

(2) INFORMATION FOR SEQ ID NO:7:

10

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

20 Cys Arg Ile Lys Gln Ile Ile Asn Arg Trp Gln Glu Val Gly Lys Ala
1 5 10 15
25 Ile Tyr Ala Pro Pro Ile Ser Gly Gln Ile Arg Cys
20 25

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Cys Arg Ile Lys Gln Ile Val Asn Met Trp Gln Arg Val Gly Gln Ala
1 5 10 15
40 Met Tyr Ala Pro Pro Ile Lys Gly Val Ile Lys Cys
20 25

(2) INFORMATION FOR SEQ ID NO:9:

45 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Cys Lys Ile Lys Gln Ile Ile Asn Met Trp Gln Gly Ala Gly Gln Ala
 1 5 10 15

Met Tyr Ala Pro Pro Ile Ser Gly Thr Ile Asn Cys
 5 20 25

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 28 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Cys Arg Ile Lys Gln Phe Ile Asn Met Trp Gln Glu Val Gly Lys Ala
 1 5 10 15

20

Met Tyr Ala Pro Pro Ile Ser Gly Gln Ile Arg Cys
 20 25

(2) INFORMATION FOR SEQ ID NO:11:

25

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

35 Cys Lys Ile Lys Gln Ile Ile Asn Met Trp Gln Lys Val Gly Lys Ala
 1 5 10 15

Met Tyr Ala Pro Pro Ile Ser Gly Gln Ile Arg Cys
 20 25

40

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 28 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Cys Arg Ile Lys Gln Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala

1 5 10
Met Tyr Ala Pro Pro Ile Ser Gly Gln Ile Arg Cys
 20 25

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 28 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Cys Lys Ile Lys Gln Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala
1 5 10 15

20 Met Tyr Ala Pro Pro Ile Glu Gly Gln Ile Arg Cys
20 25

(2) INFORMATION FOR SEQ ID NO:14:

25 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 92 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

35 Ser Gly Gly Asp Pro Glu Ile Val Met His Ser Phe Asn Cys Gly Gly
 5 10 15

Glu Phe Phe Tyr Cys Asn Thr Ser Pro Leu Phe Asn Ser Ile Trp Asn
20 25 30

40 Gly Asn Asn Thr Trp Asn Asn Thr Thr Gly Ser Asn Asn Asn Ile Thr
35 40 45

Leu Gln Cys Lys Ile Lys Gln Ile Ile Asn Met Trp Gln Lys Val Gly
50 55 60

45 Lys Ala Met Tyr Ala Pro Pro Ile Glu Gly Gln Ile Arg Cys Ser Ser
65 70 75 80

50 Asn Ile Thr Gly Leu Leu Leu Thr Arg Asp Gly Gly 85 90

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 28 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

10 Cys Lys Ile Lys Gln Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala
 1 5 10 15

Met Tyr Ala Pro Pro Ile Glu Gly Gln Ile Arg Cys
 20 25

15

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 28 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

20

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Cys Lys Ile Lys Gln Ile Ile Asn Met Trp Gln Ala Val Gly Lys Ala
 1 5 10 15

30 Met Tyr Ala Pro Pro Ile Glu Gly Gln Ile Arg Cys
 20 25

(2) INFORMATION FOR SEQ ID NO:17:

- 35 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 28 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

45 Cys Ala Ile Lys Gln Ile Ile Asn Met Trp Gln Lys Val Gly Lys Ala
 1 5 10 15

Met Tyr Ala Pro Pro Ile Glu Gly Gln Ile Arg Cys
 20 25

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

10 Cys Lys Ile Ala Gln Ile Ile Asn Met Trp Gln Lys Val Gly Lys Ala
1 5 10 15
Met Tyr Ala Pro Pro Ile Glu Gly Gln Ile Arg Cys
20 25

(25) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 28 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

25 Cys Lys Ile Lys Lys Gln Ile Ile Asn Met Trp Gln Lys Val Gly Ala Ala
1 5 10 15
30 Met Tyr Ala Pro Pro Ile Glu Gly Gln Ile Arg Cys
20 25

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 28 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

45 Cys Lys Ile Lys Lys Gln Ile Ile Asn Met Trp Gln Lys Val Gly Lys Ala
1 5 10 15
Met Tyr Ala Pro Pro Ile Ala Gly Gln Ile Arg Cys
20 25

(2) INFORMATION FOR SEQ ID NO:21:

50

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 amino acids

(B) TYPE: amino acid
(D) TOPOLOGY: linear

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Cys Arg Ile Lys Gln Phe Ile Asn Met Trp Gln Glu Val Gly Lys Ala
1 5 10 15

10

Met Tyr Ala Pro Pro Ile Ser Gly Gln Ile Arg Cys
20 25

(2) INFORMATION FOR SEQ ID NO:22:

15

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

25 Cys Lys Ile Lys Gln Phe Ile Asn Met Trp Gln Lys Val Gly Lys Ala
1 5 10 15

Met Tyr Ala Pro Pro Ile Glu Gly Gln Ile Arg Cys
20 25

30

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 amino acids

35

(B) TYPE: amino acid

(D) TOPOLOGY: linear

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Cys Lys Ile Lys Gln Phe Ile Asn Met Trp Gln Glu Val Gly Lys Ala
1 5 10 15

45 Met Tyr Ala Pro Pro Ile Glu Gly Gln Ile Arg Cys
20 25

(2) INFORMATION FOR SEQ ID NO:24:

50

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Phe Ile Asn Met Trp Gln Glu Val Gly Lys Ala Met Tyr Ala Pro Pro
1 5 10 15

10 Ile Ser

(2) INFORMATION FOR SEQ ID NO:25:

15 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Met Trp Gln Glu Val Gly Lys Ala Met Tyr Ala Pro
25 1 5 10

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 14 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Gly Lys Ala Met Tyr Ala Pro Pro Ile Lys Gly Gln Ile Arg
1 5 10
40

WHAT IS CLAIMED IS:

1. A method for making an HIV gp120 subunit vaccine for a geographic region comprising the steps of:
 - 5 a. determining a neutralizing epitope in the V2 or C4 domain of gp120 of HIV isolates from the geographic region;
 - b. selecting an HIV strain having gp120 which has a neutralizing epitope in the V2 or C4 domain which is common among isolates in the geographic region; and
 - 10 c. making an HIV gp120 subunit vaccine from the selected isolate.
2. The method of Claim 1 wherein the neutralizing epitope is determined by determining the amino acid sequence for at least a portion of the V2 or C4 domain.
3. The method of Claim 2 wherein the amino acid sequence is determined by sequencing DNA encoding at least a portion of the V2 or C4 domain.
4. The method of Claim 3 wherein a plurality of isolates having different amino acid sequences for the V2 and C4 domains are selected.
5. The method of Claim 4 wherein a plurality of isolates having different amino acid sequences for the V3 domain is selected.
6. A method for making an HIV gp120 subunit vaccine for a geographic region comprising the steps of:
 - 30 a. determining neutralizing epitopes for the V2, V3, and C4 domains of gp120 from HIV isolates from the geographic region;
 - b. selecting at least two HIV isolates having different neutralizing epitopes in the V2, V3, or C4 domain; and
 - 35 c. making an HIV gp120 subunit vaccine from the selected isolates.

7. The method of Claim 6 wherein each of the selected isolates have one of the most common neutralizing epitopes.
8. A method for making an HIV gp120 subunit vaccine for a geographic region comprising the steps of:
 - a. determining the neutralizing epitopes for HIV isolates and the percentage of HIV infections attributable to each strain present in the region;
 - 10 b. selecting at least two HIV strains which have the most common neutralizing epitopes in the V2, V3, and C4 domains in the geographic region; and
 - 15 c. making an HIV gp120 subunit vaccine from the selected isolates.
9. The method of Claim 8 wherein the isolates are primary patient isolates.
10. The method of Claim 8 wherein the geographic region is North America and the amino acid
20 sequence of gp120 from the HIV isolates MN and GNE₁ are selected.
11. The method of Claim 10 wherein the GNE₁ isolate is also selected.
12. A multivalent HIV gp120 subunit vaccine.
- 25 13. The vaccine of Claim 12 wherein gp120 present in the vaccine is from at least two HIV isolates which have a different neutralizing epitope in the V2 or C4 domain of gp120.
14. The vaccine of Claim 12 wherein gp120 present in
30 the vaccine is from at least two HIV isolates which have a different neutralizing epitope in the V3 domain of gp120.
15. The vaccine of Claim 12 wherein each isolate has a
35 or C4 domains of gp120.

16. The vaccine of Claim 1 wherein gp120 present in the vaccine is from the MN and GNE₈ strains of HIV.
17. The vaccine of Claim 1 wherein gp120 from the GNE₁₆ strain of HIV is also present in the vaccine.
- 5 18. A DNA sequence of less than 5 kilobases encoding gp120 from GNE₈ and having the nucleotide sequence illustrated in Table 1.
19. A DNA sequence of less than 5 kilobases encoding gp120 from GNE₁₆ and having the nucleotide sequence
10 illustrated in Table 2.
20. An expression construct comprising DNA encoding gp120 selected from the group consisting of GNE₈-gp120 and GNE₁₆-gp120 under the transcriptional and translational control of a heterologous
15 promoter.
21. The expression construct of Claim 20 wherein the promoter is a eukaryotic promoter.
22. The expression construct of Claim 21 wherein the DNA encoding gp120 is joined to a heterologous
20 signal sequence.
23. An isolated GNE₈-gp120 polypeptide having the amino acid sequence illustrated in Table 1.
24. An isolated GNE₁₆-gp120 polypeptide having the amino acid sequence illustrated in Table 2.
- 25 25. An improved serotyping method for HIV strains comprising determining the serotypes of the V2, V3, and C4 domains of gp120.
26. A truncated gp120 sequence which sequence is free from the C5 domain.
- 30 27. The truncated gp120 sequence of Claim 26 wherein the sequence is additionally free from the carboxy terminus through amino acid residue 453 of the gp120 V5 domain.
28. The truncated gp120 sequence of Claim 27 wherein
35 the sequence is additionally free from the gp120 V5 domain.

29. The truncated gp120 sequence of Claim 26 wherein the sequence is additionally free from the gp120 signal sequence.
- 5 30. The truncated gp120 sequence of Claim 29 wherein the sequence is additionally free from the carboxy terminus through amino acid residue 111 of the gp120 C1 domain.
- 10 31. The truncated gp120 sequence of Claim 29 wherein the sequence is additionally free from the carboxy terminus through amino acid residue 117 of the gp120 C1 domain.
- 15 32. The truncated gp120 sequence of Claim 26 wherein the sequence is free from the amino terminus of gp120 through residue 111 of the C1 domain and residue 453 through the carboxy terminus of gp120.
33. The truncated gp120 sequence of Claim 26 wherein the sequence is produced by recombinant engineering.




FIG. 1




FIG. 2




FIG. 3A




FIG. 3B

3 / 10

418	445			
CKIKQIINMWQKGKAMYAPPIEGQIRC	MN _{GNE}	(SEQ. ID. NO. 3)		
-----E-----	MN ₁₉₈₄	(SEQ. ID. NO. 4)		
-R-----E-----K-----	JRCSF	(SEQ. ID. NO. 5)		
-R-----E-----N-----	Z6	(SEQ. ID. NO. 6)		
-R-----R-E-I-----S-----	NY5	(SEQ. ID. NO. 7)		
-R-----V-----K-V-K-----	Z321	(SEQ. ID. NO. 8)		
-----GA-Q-----S-T-N-----	A244	(SEQ. ID. NO. 9)		
-R-F-----E-----S-----	LAI _{IIIB}	LAI _{BRU} , LAI _{HXB3} (SEQ. ID. NO. 10)		
-I-----K-----S-----	LAI _{HXB2}	(SEQ. ID. NO. 11)		
-R-I-----E-----S-----	LAI _{BH10}	LAI _{HXB3} (SEQ. ID. NO. 12)		
-----E-----S-----	MN ₁₉₈₄	(SEQ. ID. NO. 13)		

FIG. 4

418	445		
CKIKQIINMWQKVGKAMYAPPIEGQIRC	MNGNE	(SEQ. ID. NO. 3)	
-----E-----	MN. 429E	(SEQ. ID. NO. 15)	
-----A-----	MN. 429A	(SEQ. ID. NO. 16)	
-A-----	MN. 419A	(SEQ. ID. NO. 17)	
-----A-----	MN. 421A	(SEQ. ID. NO. 18)	
-----A-----	MN. 432A	(SEQ. ID. NO. 19)	
-----A-----	MN. 440A	(SEQ. ID. NO. 20)	
-R-F-E-----S-----	LAI IIB	(SEQ. ID. NO. 21)	
-----F-----	MN. 423F	(SEQ. ID. NO. 22)	
-----F-----	MN. 423F, 429E	(SEQ. ID. NO. 23)	

FIG. 5








FIG. 6



8/10




FIG. 8

8 / 10




FIG. 9




FIG. 10

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/06036

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : IPC(5): A61K 39/12, 37/02; C12N 15/00; C07K 3/00; C07H 15/12

US CL : 424/89; 435/172.1, 320.1; 530/333, 350; 536/27

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/89; 435/172.1, 320.1; 530/333, 350; 536/27

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, Dialog, Search terms: HIV, vaccine, variable domains, constant domains, envelope, neutralizing epitopes

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Septieme Colloque Cent Gardes, issued 1992, M. Klein, "Immunogenicity of Synthetic HIV-1 T-B Tandem Epitopes", pages 169-174, see entire article.	5-12, 14, 28-33
Y	Journal of Virology, Vol. 66, No. 9, issued September 1992, M. Thali et al, "Discontinuous, Conserved Neutralization Epitopes Overlapping the CD4-Binding Region of Human Immunodeficiency Virus Type 1 gp120 Envelope Glycoprotein", pages 5635-5641, see entire article.	1-11



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z

document member of the same patent family

Date of the actual completion of the international search

30 SEPTEMBER 1994

Date of mailing of the international search report

20 OCT 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

LYNETTE F. SMITH

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/06036

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	AIDS Research and Human Retroviruses, Vol. 5, No. 1, issued 1989, "Patterns of Antibody Recognition of Selected Conserved Amino Acid Sequences from the HIV Envelope in Sera from Different Stages of HIV Infection, pages 33-39, see entire article.	1-4
Y	WO, A, 91/ 15512 (GREGORY ET AL.) 17 October 1991, see entire publication.	1-33
Y	Proceedings of the National Academy of Sciences USA, Vol. 89, issued January 1992, Broliden et al, "Identification of Human Neutralization-Inducing Regions of the Human Immunodeficiency Virus Type 1 Envelope Glycoproteins", pages 461-465, see entire article.	5-12, 14, 28-33
Y	Proceedings of the National Academy of Sciences USA, Vol. 89, issued September 1989, Javaherian et al, "Principal Neutralizing Domain of the Human Immunodeficiency Virus Type 1 Envelope Protein", pages 6768-6772, see entire article.	5-12, 14, 28-33